



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/385, 39/395, G01N 33/53, C12P 21/06, 21/04, C07K 16/00		A1	(11) International Publication Number: WO 98/48837 (43) International Publication Date: 5 November 1998 (05.11.98)
(21) International Application Number: PCT/US98/08654 (22) International Filing Date: 30 April 1998 (30.04.98) (30) Priority Data: 60/044,449 30 April 1997 (30.04.97) US 60/050,472 23 June 1997 (23.06.97) US 60/063,074 27 October 1997 (27.10.97) US 60/067,341 2 December 1997 (02.12.97) US (71) Applicant: ENZON, INC. [US/US]; 20 Kingsbridge Road, Piscataway, NJ 08854-3963 (US). (72) Inventors: WHITLOW, Marc; 3675 May Road, El Sobrante, CA 94803 (US). SHORR, Robert, G., L.; 28 Brookfall Road, Edison, NJ 08817 (US). FILPULA, David, R.; 269 Carlton Club Drive, Piscataway, NJ 08854 (US). LEE, Lihsyng, S.; 22 Van Wyck Drive, Princeton Junction, NJ 08550 (US). (74) Agents: GOLDSTEIN, Jorge, A. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: POLYALKYLENE OXIDE-MODIFIED SINGLE CHAIN POLYPEPTIDES			
(57) Abstract			
<p>The present invention relates to the chemical modification of single chain polypeptides by means of covalent attachment of strands of poly(ethylene glycol) PEG and similar poly(alkylene oxides) to single chain polypeptide binding molecules that have the three dimensional folding and, thus, the binding ability and specificity, of the variable region of an antibody. Such preparations of modified single chain polypeptide binding molecules have reduced immunogenicity and antigenicity as well as having a longer halflife in the bloodstream as compared to the parent polypeptide. These beneficial properties of the modified single chain polypeptide binding molecules make them very useful in a variety of therapeutic applications. The invention also relates to multivalent antigen-binding molecules capable of PEGylation. Compositions of, genetic constructions for, methods of use, and methods for producing PEGylated antigen-binding proteins are disclosed.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Polyalkylene Oxide-Modified Single Chain Polypeptides

Background of the Invention

1. Field of the Invention

5 The present invention relates to the chemical modification of single chain polypeptides by means of covalent attachment of strands of poly(ethylene glycol) PEG and similar poly(alkylene oxides) to single chain polypeptide binding molecules that have the three dimensional folding and, thus, the binding ability and specificity, of the variable region of an antibody. Such preparations of modified
10 single chain polypeptide binding molecules have reduced immugenicity and antigenicity as well as having a longer halflife in the bloodstream as compared to the parent polypeptide. These beneficial properties of the modified single chain polypeptide binding molecules make them very useful in a variety of therapeutic applications. The invention also relates to multivalent antigen-binding molecules
15 capable of PEGylation. Compositions of, genetic constructions for, methods of use, and methods for producing PEGylated antigen-binding proteins are disclosed.

2. Description of Related Art

 Antibodies are proteins generated by the immune system to provide a specific molecule capable of complexing with an invading molecule, termed an
20 antigen. Natural antibodies have two identical antigen-binding sites, both of which are specific to a particular antigen. The antibody molecule "recognizes" the antigen by complexing its antigen-binding sites with areas of the antigen termed

-2-

epitopes. The epitopes fit into the conformational architecture of the antigen-binding sites of the antibody, enabling the antibody to bind to the antigen.

The antibody molecule is composed of two identical heavy and two identical light polypeptide chains, held together by interchain disulfide bonds. The remainder of this discussion on antibodies will refer only to one pair of light/heavy chains, as each light/heavy pair is identical. Each individual light and heavy chain folds into regions of approximately 110 amino acids, assuming a conserved three-dimensional conformation. The light chain comprises one variable region (V_L) and one constant region (C_L), while the heavy chain comprises one variable region (V_H) and three constant regions (C_{H1} , C_{H2} and C_{H3}). Pairs of regions associate to form discrete structures. In particular, the light and heavy chain variable regions associate to form an "Fv" area which contains the antigen-binding site. The constant regions are not necessary for antigen binding and in some cases can be separated from the antibody molecule by proteolysis, yielding biologically active (i.e., binding) variable regions composed of half of a light chain and one quarter of a heavy chain.

Further, all antibodies of a certain class and their F_{ab} fragments (i.e., fragments composed of V_L , C_L , V_H , and C_{H1}) whose structures have been determined by x-ray crystallography show similar variable region structures despite large differences in the sequence of hypervariable segments even when from different animal species. The immunoglobulin variable region seems to be tolerant towards mutations in the antigen-binding loops. Therefore, other than in the hypervariable regions, most of the so-called "variable" regions of antibodies, which are defined by both heavy and light chains, are, in fact, quite constant in their three dimensional arrangement. See for example, Huber, R., *Science* 233:702-703 (1986)).

Recent advances in immunobiology, recombinant DNA technology, and computer science have allowed the creation of *single* polypeptide chain molecules that bind antigen. These single-chain antigen-binding molecules ("SCA") or single-chain variable fragments of antibodies ("sFv") incorporate a linker

polypeptide to bridge the individual variable regions, V_L and V_H , into a single polypeptide chain. A description of the theory and production of single-chain antigen-binding proteins is found in Ladner *et al.*, U.S. Patent Nos. 4,946,778, 5,260,203, 5,455,030 and 5,518,889. The single-chain antigen-binding proteins produced under the process recited in the above U.S. patents have binding specificity and affinity substantially similar to that of the corresponding Fab fragment. A computer-assisted method for linker design is described more particularly in Ladner *et al.*, U.S. Patent Nos. 4,704,692 and 4,881,175, and WO 94/12520.

The *in vivo* properties of sFv (SCA) polypeptides are different from MAbs and antibody fragments. Due to their small size, sFv (SCA) polypeptides clear more rapidly from the blood and penetrate more rapidly into tissues (Milenic, D.E. *et al.*, *Cancer Research* 51:6363-6371 (1991); Colcher *et al.*, *J. Natl. Cancer Inst.* 82:1191 (1990); Yokota *et al.*, *Cancer Research* 52:3402 (1992)). Due to lack of constant regions, sFv (SCA) polypeptides are not retained in tissues such as the liver and kidneys. Due to the rapid clearance and lack of constant regions, sFv (SCA) polypeptides will have low immunogenicity. Thus, sFv (SCA) polypeptides have applications in cancer diagnosis and therapy, where rapid tissue penetration and clearance, and ease of microbial production are advantageous.

A multivalent antigen-binding protein has more than one antigen-binding site. A multivalent antigen-binding protein comprises two or more single-chain protein molecules. Enhanced binding activity, di- and multi-specific binding, and other novel uses of multivalent antigen-binding proteins have been demonstrated. See, Whitlow, M., *et al.*, *Protein Engng.* 7:1017-1026 (1994); Hoogenboom, H.R., *Nature Biotech.* 15:125-126 (1997); and WO 93/11161.

Ladner *et al.* also discloses the use of the single chain antigen binding molecules in diagnostics, therapeutics, *in vivo* and *in vitro* imaging, purifications, and biosensors. The use of the single chain antigen binding molecules in immobilized form, or in detectably labeled forms is also disclosed, as well as conjugates of the single chain antigen binding molecules with therapeutic agents,

such as drugs or specific toxins, for delivery to a specific site in an animal, such as a human patient.

Whitlow *et al.* (*Methods: A Companion to Methods in Enzymology* 2(2):97-105 (June, 1991)) provide a good review of the art of single chain antigen binding molecules and describe a process for making them.

In U.S. Patent 5,091,513, Huston *et al.* discloses a family of synthetic proteins having affinity for preselected antigens. The contents of U.S. Patent 5,091,513 are incorporated by reference herein. The proteins are characterized by one or more sequences of amino acids constituting a region that behaves as a biosynthetic antibody binding site (BABS). The sites comprise (1) noncovalently associated or disulfide bonded synthetic V_H and V_L regions, (2) V_H - V_L or V_L - V_H single chains wherein the V_H and V_L are attached to a polypeptide linker, or (3) individual V_H or V_L domains. The binding domains comprises complementarity determining regions (CDRs) linked to framework regions (FRs), which may be derived from separate immunoglobulins.

U.S. Patent 5,091,513 also discloses that three subregions (the CDRs) of the variable domain of each of the heavy and light chains of native immunoglobulin molecules collectively are responsible for antigen recognition and binding. These CDRs consist of one of the hypervariable regions or loops and of selected amino acids or amino acid sequences disposed in the framework regions that flank that particular hypervariable region. It is said that framework regions from diverse species are effective in maintaining CDRs from diverse other species in proper conformation so as to achieve true immunochemical binding properties in a biosynthetic protein.

U.S. Patent 5,091,513 includes a description of a chimeric polypeptide that is a single chain composite polypeptide comprising a complete antibody binding site. This single chain composite polypeptide is described as having a structure patterned after tandem V_H and V_L domains, with a carboxyl terminal of one attached through an amino acid sequence to the amino terminal of the other. It thus comprises an amino acid sequence that is homologous to a portion of the

variable region of an immunoglobulin heavy chain (V_H) peptide bonded to a second amino acid sequence that was homologous to a portion of the variable region of an immunoglobulin light chain (V_L).

5 The covalent attachment of strands of a polyalkylene glycol to a polypeptide molecule is disclosed in U.S. Patent No. 4,179,337 to Davis *et al.*, as well as in Abuchowski and Davis "Enzymes as Drugs," Holcenberg and Roberts, Eds., pp. 367-383, John Wiley and Sons, New York (1981). These references disclosed that proteins and enzymes modified with polyethylene glycols have reduced immunogenicity and antigenicity and have longer lifetimes in the
10 bloodstream, compared to the parent compounds. The resultant beneficial properties of the chemically modified conjugates are very useful in a variety of therapeutic applications.

Although amino acid sequences such as the single chain polypeptides described above, and fusion proteins thereof, have not been associated with
15 significant antigenicity in mammals, it has been desirable to prolong the circulating life and even further reduce the possibility of an antigenic response. The relatively small size of the polypeptides and their delicate structure/activity relationship, however, have made polyethylene glycol modification difficult and unpredictable. Most importantly, it was unknown how to modulate retained activity of the
20 polypeptides after conjugation with polymers, such as PEG.

To effect covalent attachment of polyethylene glycol (PEG) or polyalkylene oxides to a protein, the hydroxyl end groups of the polymer must first be converted into reactive functional groups. This process is frequently referred to as "activation" and the product is called "activated PEG" or activated
25 polyalkylene oxide. Methoxy poly(ethylene glycol) (mPEG), capped on one end with a functional group, reactive towards amines on a protein molecule, is used in most cases.

The activated polymers are reacted with a therapeutic agent having nucleophilic functional groups that serve as attachment sites. One nucleophilic
30 functional group commonly used as an attachment site is the ϵ -amino groups of

lysines. Free carboxylic acid groups, suitably activated carbonyl groups, oxidized carbohydrate moieties and mercapto groups have also been used as attachment sites.

5 The hydroxyl group of PEG has been activated with cyanuric chloride and the resulting compound is then coupled with proteins (Abuchowski *et al.*, *J. Biol. Chem.* 252:3578 (1977); Abuchowski & Davis, *supra* (1981)). However, there are disadvantages in using this method, such as the toxicity of cyanuric chloride and its non-specific reactivity for proteins having functional groups other than amines, such as free essential cysteine or tyrosine residues.

10 In order to overcome these and other disadvantages, alternative activated PEGs, such as succinimidyl succinate derivatives of PEG ("SS-PEG"), have been introduced (Abuchowski *et al.*, *Cancer Biochem. Biophys.* 7:175-186 (1984)). SS-PEG reacts quickly with proteins (30 minutes) under mild conditions yielding active yet extensively modified conjugates.

15 Zalipsky, in U.S. Patent No. 5,122,614, discloses poly(ethylene glycol)-N-succinimide carbonate and its preparation. This form of the polymer is said to react readily with the amino groups of proteins, as well as low molecular weight peptides and other materials that contain free amino groups.

20 Other linkages between the amino groups of the protein, and the PEG are also known in the art, such as urethane linkages (Veronese *et al.*, *Appl. Biochem. Biotechnol.* 11:141-152 (1985)), carbamate linkages (Beauchamp *et al.*, *Analyt. Biochem.* 131:25-33 (1983)), and others.

25 Suzuki *et al.* (*Biochimica et Biophysica Acta*, 788: 248-255 (1984)) covalently couples immunoglobulin G (IgG) to poly(ethylene glycol) that has previously been activated by cyanuric chloride. The coupled IgG was studied for physicochemical and biological properties such as molecular structure, size-exclusion chromatographic behavior, surface activity, interfacial aggregability, heat aggregability inducing nonspecific complement activation, and antigen-binding activity. The poly(ethylene glycol) coupling to IgG increased the apparent
30 Stokes' radius and the surface activity of IgG and stabilized IgG on heating and/or

on exposure to interfaces, while no structural denaturation of IgG was observed. The suppressed nonspecific aggregability was interpreted mainly by difficulty in association between the modified IgG molecules. These results indicated the use of the poly(ethylene glycol)-coupled IgG as an intravenous preparation and also
5 as an additive stabilizing intact IgG for intravenous use.

Sharp *et al.* (*Analytical Biochemistry* 154: 110-117 (1986)) investigated the possibility of producing biospecific affinity ligands for separating cells in two polymer aqueous phase systems on the basis of cell surface antigens. Rabbit anti-human erythrocyte IgG was reacted with cyanuric chloride-activated monomethyl
10 poly(ethylene glycol) fractions (molecular weights approximately 200, 1900, and 5000) at various molar ratios of PEG to protein lysine groups. The partition coefficient of the protein in a Dextran/PEG two phase system increased with increasing degree of modification and increasing PEG molecular weight. There was a concomitant loss in ability to agglutinate human erythrocytes.

Tullis, in U.S. Patent No. 4,904,582, describes oligonucleotide conjugates wherein the oligonucleotides are joined through a linking arm to a hydrophobic moiety, which could be a polyalkyleneoxy group. The resulting conjugates are said to be more efficient in membrane transport, so as to be capable of crossing the membrane and effectively modulating a transcriptional system. In this way, the
15 compositions can be used in vitro and in vivo, for studying cellular processes, protecting mammalian hosts from pathogens, and the like.

Excessive polymer conjugation and/or conjugation involving a therapeutic moiety's active site where groups associated with bioactivity are found, however, often result in loss of activity and, thus, therapeutic usefulness. This is often the
25 case with lower molecular weight peptides which have few attachment sites not associated with bioactivity. For example, Benhar *et al.* (*Bioconjugate Chem.* 5:321-326 (1994)) observed that PEGylation of a recombinant single-chain immunotoxin resulted in the loss of specific target immunoreactivity of the immunotoxin. The loss of activity of the immunotoxin was the result of PEG
30 conjugation at two lysine residues within the antibody-combining region of the

immunotoxin. To overcome this problem, Benhar *et al.* replaced these two lysine residues with arginine residues and were able to obtain an active immunotoxin that was 3-fold more resistant to inactivation by derivatization.

Another suggestion for overcoming these problems discussed above is to use longer, higher molecular weight polymers. These materials, however, are difficult to prepare and expensive to use. Further, they provide little improvement over more readily available polymers.

Another alternative suggested is to attach two strands of polymer via a triazine ring to amino groups of a protein. See, for example, *Enzyme* 26:49-53 (1981) and *Proc. Soc. Exper. Biol. Med.*, 188:364-369 (1988). However, triazine is a toxic substance that is difficult to reduce to acceptable levels after conjugation. Thus, non-triazine-based activated polymers would offer substantial benefits to the art.

Summary of the Invention

The present invention relates to polyalkylene oxide/amino acid sequence conjugates and processes for preparing them. Suitable amino acid sequences are peptides, such as, single chain polypeptides having binding affinity for an antigen, for example, those described by Ladner *et al.* in U.S. Patent No. 4,946,778 and Huston *et al.* in U.S. Patent No. 5,091,513.

More particularly, the present invention relates to a physiologically active, substantially non-immunogenic polypeptide conjugate containing at least one polyalkylene oxide strand coupled to a single chain polypeptide having binding affinity for an antigen. The single chain polypeptide includes:

- (a) a first polypeptide comprising the binding portion of the light chain variable region of an antibody;
- (b) a second polypeptide comprising the binding portion of the heavy chain variable region of an antibody; and

- (c) at least one peptide linker linking said first and second polypeptides (a) and (b) into said single chain polypeptide having binding affinity for the antigen.

In another aspect, the present invention relates to a process for preparing physiologically active, substantially non-immunogenic polypeptide compositions. The process includes coupling a polyalkylene oxide to a single chain polypeptide having the attributes described above. Preferably, the poly(alkylene oxides) used herein are poly(ethylene glycols) that have been activated for coupling to the target polypeptide.

The invention is also directed to a single-chain antigen-binding polypeptide-polyalkylene oxide conjugate, comprising:

- (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;
 - (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and
 - (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,
- wherein the single-chain antigen-binding polypeptide-polyalkylene oxide conjugate has an antigen binding affinity within a range of about one-fold to about ten-fold of the antigen binding affinity of the native, unconjugated form of the single-chain antigen-binding polypeptide.

The invention is also directed to a single-chain antigen-binding polypeptide-polyalkylene oxide conjugate, comprising:

- (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;
- (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and
- (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,

-10-

wherein the single-chain antigen-binding polypeptide-polyalkylene oxide conjugate has an antigen binding affinity within about ten-fold of the antigen binding affinity of the native, unconjugated form of the single-chain antigen-binding polypeptide.

The invention is also directed to a single-chain antigen-binding polypeptide
5 -polyalkylene oxide conjugate, comprising:

- (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;
 - (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and
 - 10 (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,
- wherein the single-chain antigen-binding polypeptide-polyalkylene oxide conjugate has an antigen binding affinity within about five-fold of the antigen binding affinity of the native, unconjugated form of the single-chain antigen-binding polypeptide.

15 The invention is also directed to a single-chain antigen-binding polypeptide -polyalkylene oxide conjugate, comprising:

- (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;
 - (b) a second polypeptide comprising the antigen binding portion of the
20 variable region of an antibody heavy or light chain; and
 - (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,
- wherein the single-chain antigen-binding polypeptide-polyalkylene oxide conjugate has an antigen binding affinity within about two-fold of the antigen binding affinity
25 of the native, unconjugated form of the single-chain antigen-binding polypeptide.

The invention is also directed to a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

- (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

-11-

(b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,

5 wherein the single-chain antigen-binding polypeptide has at least one Cys residue wherein the Cys residue is capable of polyalkylene oxide conjugation and the Cys residue is located at a position selected from the group consisting of (i) the amino acid position 11, 12, 13, 14 or 15 of the light chain variable region; (ii) the amino acid position 77, 78 or 79 of the light chain variable region; (iii) the amino acid
10 position 11, 12, 13, 14 or 15 of the heavy chain variable region; (iv) the amino acid position 82B, 82C or 83 of the heavy chain variable region; (v) any amino acid position of the peptide linker; (vi) adjacent to the C-terminus of polypeptide (a) or (b); and (vii) combinations thereof, wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an
15 antigen.

The invention is also directed to a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

(a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

20 (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,

wherein the single-chain antigen-binding polypeptide has at least three consecutive
25 Lys residues wherein the consecutive Lys residues are capable of polyalkylene oxide conjugation and any one of the consecutive Lys residues is located at a position selected from the group consisting of (i) any amino acid position of the peptide linker; (ii) adjacent to the C-terminus of polypeptide (a) or (b); and (iii) combinations thereof, wherein the polyalkylene oxide conjugated single-chain
30 antigen-binding polypeptide is capable of binding an antigen. These consecutive

lysine residues in the sFv (SCA) protein (i.e., oligo-lysine sFv) generate a "hot spot" for polyalkylene oxide conjugation.

The invention is also directed to a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

- 5 (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;
- (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and
- (c) a peptide linker linking the first and second polypeptides (a) and
- 10 (b) into a single chain polypeptide having an antigen binding site, wherein the single-chain antigen-binding polypeptide has at least two consecutive Cys residue wherein the consecutive Cys residues are capable of polyalkylene oxide conjugation and any one of the consecutive Cys residues is located at a position selected from the group consisting of (i) any amino acid position of the peptide linker; (ii) adjacent to the C-terminus of polypeptide (a) or (b); and (iii)
- 15 combinations thereof, wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen. These consecutive cysteine residues in the sFv (SCA) protein (i.e., oligo-cysteine sFv) generate a "hot spot" for polyalkylene oxide conjugation.

- 20 The invention is further directed to a genetic sequence encoding a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

- (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;
- 25 (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and
- (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site, wherein the single-chain antigen-binding polypeptide has at least one Cys residue
- 30 wherein the Cys residue is capable of polyalkylene oxide conjugation and the Cys

residue is located at a position selected from the group consisting of (i) the amino acid position 11, 12, 13, 14 or 15 of the light chain variable region; (ii) the amino acid position 77, 78 or 79 of the light chain variable region; (iii) the amino acid position 11, 12, 13, 14 or 15 of the heavy chain variable region; (iv) the amino acid position 82B, 82C or 83 of the heavy chain variable region; (v) any amino acid position of the peptide linker; (vi) adjacent to the C-terminus of polypeptide (a) or (b); and (vii) combinations thereof, wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

10 The invention is further directed to a genetic sequence encoding a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

 (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

15 (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

 (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,

wherein the single-chain antigen-binding polypeptide has at least three consecutive

20 Lys residue wherein the consecutive Lys residues are capable of polyalkylene oxide conjugation and any one of the consecutive Lys residues is located at a position selected from the group consisting of (i) any amino acid position of the peptide linker; (ii) adjacent to the C-terminus of polypeptide (a) or (b); and (iii) combinations thereof, wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen. These consecutive lysine residues in the sFv (SCA) protein (i.e., oligo-lysine sFv) generate a "hot spot" for polyalkylene oxide conjugation.

25 The invention is further directed to a genetic sequence encoding a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

30

-14-

(a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

5 (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site, wherein the single-chain antigen-binding polypeptide has at least two consecutive Cys residue wherein the consecutive Cys residues are capable of polyalkylene oxide conjugation and any one of the consecutive Cys residues is located at a
10 position selected from the group consisting of (i) any amino acid position of the peptide linker; (ii) adjacent to the C-terminus of polypeptide (a) or (b); and (iii) combinations thereof, wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen. These consecutive cysteine residues in the sFv (SCA) protein (i.e., oligo-cysteine sFv) generate a
15 "hot spot" for polyalkylene oxide conjugation.

The genetic sequence may be DNA or RNA.

The invention is directed to a replicable cloning or expression vehicle comprising the above described DNA sequence. The invention is also directed to such vehicle which is a plasmid. The invention is further directed to a host cell
20 transformed with the above described DNA. The host cell may be a bacterial cell, a yeast cell or other fungal cell, an insect cell or a mammalian cell line. A preferred host is *Pichia pastoris*.

The invention is directed to a method of producing a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

25 (a) providing a first genetic sequence encoding a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) providing a second genetic sequence encoding a second polypeptide comprising the antigen binding portion of the variable region of an
30 antibody heavy or light chain; and

- (c) linking the first and second genetic sequences (a) and (b) with a third genetic sequence encoding a peptide linker into a fourth genetic sequence encoding a single chain polypeptide having an antigen binding site, wherein the single-chain antigen-binding polypeptide has at least one Cys residue wherein the Cys residue is capable of polyalkylene oxide conjugation and the Cys residue is located at a position selected from the group consisting of (i) the amino acid position 11, 12, 13, 14 or 15 of the light chain variable region; (ii) the amino acid position 77, 78 or 79 of the light chain variable region; (iii) the amino acid position 11, 12, 13, 14 or 15 of the heavy chain variable region; (iv) the amino acid position 82B, 82C or 83 of the heavy chain variable region; (v) any amino acid position of the peptide linker; (vi) adjacent to the C-terminus of polypeptide (a) or (b); and (vii) combinations thereof, wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen;
- (d) transforming a host cell with the fourth genetic sequence encoding a single-chain antigen-binding polypeptide of (c); and
- (e) expressing the single-chain antigen-binding polypeptide of (c) in the host, thereby producing a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation.
- The invention is further directed to a multivalent single-chain antigen-binding protein, comprising two or more single-chain antigen-binding polypeptides, each single-chain antigen-binding polypeptide comprising:
- (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;
- (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and
- (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site, wherein the single-chain antigen-binding polypeptide has at least one Cys residue wherein the Cys residue is capable of polyalkylene oxide conjugation and the Cys

residue is located at a position selected from the group consisting of (i) the amino acid position 11, 12, 13, 14 or 15 of the light chain variable region; (ii) the amino acid position 77, 78 or 79 of the light chain variable region; (iii) the amino acid position 11, 12, 13, 14 or 15 of the heavy chain variable region; (iv) the amino acid position 82B, 82C or 83 of the heavy chain variable region; (v) any amino acid position of the peptide linker; (vi) adjacent to the C-terminus of polypeptide (a) or (b); and (vii) combinations thereof, wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

10 In the above described embodiments of the invention, the Cys polyalkylene oxide conjugation sequence may be capable of attaching a polyalkylene oxide moiety and the Cys residue is located at a position selected from the group consisting of (i') the amino acid position 77 of the light chain variable region; (ii') the amino acid position 82B of the heavy chain variable region; (iii') the amino acid position 3 of the peptide linker; (iv') adjacent to the C-terminus of polypeptide (a) or (b); (v') N-terminus and C-terminus; and (vi') combinations thereof, wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

20 In the above described embodiments of the invention, the oligo-Lys polyalkylene oxide conjugation sequence may be capable of attaching a polyalkylene oxide moiety at the oligo-Lys residues located adjacent to the C-terminus of the protein, wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

25 In the above described embodiments of the invention, the C-terminus of the second polypeptide (b) may be the native C-terminus. The C-terminus of the second polypeptide (b) may comprise a deletion of one or plurality of amino acid residue(s), such that the remaining N-terminus amino acid residues of the second polypeptide are sufficient for the polyalkylene oxide conjugated polypeptide to be capable of binding an antigen. The C-terminus of the second polypeptide may

comprise an addition of one or plurality of amino acid residue(s), such that the polyalkylene oxide conjugated polypeptide is capable of binding an antigen.

In a preferred embodiment of the invention, the first polypeptide (a) may comprise the antigen binding portion of the variable region of an antibody light
5 chain and the second polypeptide (b) comprises the antigen binding portion of the variable region of an antibody heavy chain.

The invention is also directed to a method of detecting an antigen suspected of being in a sample, comprising:

(a) contacting the sample with the polyalkylene oxide conjugated
10 polypeptide or protein of the invention, wherein the polyalkylene oxide conjugated polypeptide is conjugated to one or plurality of detectable label molecule(s), or conjugated to a carrier having one or plurality of detectable label molecule(s) bound to the carrier; and

(b) detecting whether the polyalkylene oxide conjugated single-chain
15 antigen-binding polypeptide has bound to the antigen.

The invention is further directed to a method of imaging the internal structure of an animal, comprising administering to the animal an effective amount of the polyalkylene oxide conjugated polypeptide or protein of the invention, wherein the polyalkylene oxide conjugated polypeptide is conjugated to one or
20 plurality of detectable label or chelator molecule(s), or conjugated to a carrier having one or plurality of detectable label or chelator molecule(s) bound to the carrier, and measuring detectable radiation associated with the animal. Animal includes human and nonhuman.

The invention is also directed to a method for treating a targeted disease,
25 comprising administering an effective amount of a composition comprising the polyalkylene oxide conjugated polypeptide or protein of the invention and a pharmaceutically acceptable carrier vehicle, wherein the polyalkylene oxide conjugated polypeptide is conjugated to one or plurality of bioactive molecules, such as peptides, lipids, nucleic acids (i.e., phosphate-lysine complexes), drug,
30 toxin, boron addend or radioisotope molecule(s), or conjugated to a carrier having

one or plurality of peptides, lipids, nucleic acids (i.e., phosphate-lysine complexes), drug, toxin, boron addend or radioisotope molecule(s) bound to the carrier.

Brief Description of the Drawings

5 Figure 1 is a graphical representation of three competition ELISA's in which unlabeled PEG modified CC49/212 SCA (closed squares), CC49/212 SCA (open squares), CC49 IgG (open circles), and MOPC-21 IgG (+) competed against a CC49 IgG radiolabeled with ¹²⁵I for binding to the TAG-72 antigen on a human breast carcinoma extract.

10 Figure 2 shows the DNA and protein sequence of CC49/218 SCA which has four engineered cysteine residues at the positions indicated by the codons underlined and marked by an asterisk. Also highlighted are the CDR sequences (double underlined) and the 218 linker (underlined and labeled). In addition, there are four natural cysteine residues in the protein which are involved in two disulfide
15 bonds. These are not underlined. The four engineered cysteine residues occur independently in four different mutants currently, but may be combined in the exact four-mutant codon version shown in this figure.

 Figure 3 shows the DNA and protein sequence of CC49/218 SCA with an engineered oligo-lysine C-terminal tail segment. The eight new lysine residues
20 were genetically engineered at a BstEII site and are shown underlined and marked with asterisks. Also highlighted are the CDR sequences (double underlined), the 218 linker (underlined and labeled) and selected restriction sites.

 Figure 4 is a graphical representation of three competition ELISA's in which unlabeled SC-PEG unreacted CC49/218 SCA (closed squares), CC49/218
25 SCA (open squares), unlabeled XUS-PEG unreacted CC49/218 SCA (open circles), SC-PEG modified CC49/218 SCA (closed circles), XUS-PEG modified CC49/218 SCA (open triangles), CC49 IgG (closed triangles), an Anti-FITC SCA (dashed line) or BL-3 IgG (dotted line) were competed against a CC49 IgG

radiolabeled with ^{125}I for binding to the TAG-72 antigen on a human breast carcinoma extract.

Figure 5 shows the pharmacokinetics of plasma retention of SCA and PEG-SCA. The details of the experiment are described in Example 13.

5 Figure 6 shows an SDS-PAGE of the purified CC49-multimers cross-linked by PEG5000 under reducing conditions. The details of the experiment are described in Example 14. The lanes of the gel contain the following: 1) trimeric form; 2) dimeric form; 3) dimeric form; 4) mixed population; 5) native CC49; 6) PEG-CC49 monomer; 7) PEG-CC49 monomer; 8) empty; 9) empty; and 10) molecular weight standards.

10 Figure 7 shows the binding kinetics of Mono-, Di-, Tri-, -PEG-CC49. The details of the experiment are described in Example 14. Native CC49 is represented by the solid box. PEG-mono-CC49 is represented by the open box. PEG-Di-CC49 is represented by the solid diamond. PEG-Tri-CC49 is represented
15 by the open diamond.

Figure 8 shows the results of the competition assay performed in Example 16. Nat is native CC49-SCA, C2 is PEG SC2000-CC49-SCA; GC is glyco-CC49-SCA; B* is the biotinylated CC49-SCA; C12 is the PEG-SC12,000-CC49-SCA; F5 is PEG-Flan-5000-CC49-SCA; and C20 is PEG-SC20000-CC49-SCA.

20

Description of the Embodiments

The present invention is directed to the novel combination of a polyalkylene glycol and a single chain polypeptide having binding affinity for an antigen, the polyalkylene glycol and polypeptide preferably being joined together by means of a coupling agent.

Single Chain Polypeptides

The invention relates to the discovery that polyalkylene oxide conjugated single-chain antigen-binding proteins ("SCA") or single-chain variable fragments of antibodies ("sFv"), such as PEGylated SCA proteins, have significant utility beyond that of the nonPEGylated single-chain antigen-binding proteins. In addition to maintaining an antigen binding site, a PEGylated SCA protein has a PEG moiety which reduces antigenicity and increases the half life of the modified polypeptide in the bloodstream. Accordingly, the invention is directed to monovalent and multivalent SCA proteins capable of PEGylation, compositions of monovalent and multivalent PEGylated SCA proteins, methods of making and purifying monovalent and multivalent PEGylated SCA proteins, and uses for PEGylated SCA proteins. The invention is also directed to PEGylated SCA proteins having a diagnostic or therapeutic agent covalently attached to an Cys-linked PEGylated polypeptide or an oligo-Lys linked PEGylated polypeptide.

The terms "single-chain antigen-binding molecule" (SCA) or "single-chain Fv" (sFv) are used interchangeably. They are structurally defined as comprising the binding portion of a first polypeptide from the variable region of an antibody V_L (or V_H), associated with the binding portion of a second polypeptide from the variable region of an antibody V_H (or V_L), the two polypeptides being joined by a peptide linker linking the first and second polypeptides into a single polypeptide chain, such that the first polypeptide is N-terminal to the linker and second polypeptide is C-terminal to the first polypeptide and linker. The single polypeptide chain thus comprises a pair of variable regions connected by a polypeptide linker. The regions may associate to form a functional antigen-binding site, as in the case wherein the regions comprise a light-chain and a heavy-chain variable region pair with appropriately paired complementarity determining regions (CDRs). In this case, the single-chain protein is referred to as a "single-chain antigen-binding protein" or "single-chain antigen-binding molecule."

Single-chain Fvs can and have been constructed in several ways. Either V_L is the N-terminal domain followed by the linker and V_H (a V_L -Linker- V_H construction) or V_H is the N-terminal domain followed by the linker and V_L (V_H -Linker- V_L construction). The preferred embodiment contains V_L in the N-terminal domain (*see*, Anand, N.N., *et al.*, *J. Biol. Chem.* 266:21874-21879 (1991)). Alternatively, multiple linkers have also been used. Several types of sFv (SCA) proteins have been successfully constructed and purified, and have shown binding affinities and specificities similar to the antibodies from which they were derived.

A description of the theory and production of single-chain antigen-binding proteins is found in Ladner *et al.*, U.S. Patent Nos. 4,946,778, 5,260,203, 5,455,030 and 5,518,889, and in Huston *et al.*, U.S. Patent No. 5,091,513 ("biosynthetic antibody binding sites" (BABS)), all incorporated herein by reference. The single-chain antigen-binding proteins produced under the process recited in the above patents have binding specificity and affinity substantially similar to that of the corresponding Fab fragment.

Typically, the Fv domains have been selected from the group of monoclonal antibodies known by their abbreviations in the literature as 26-10, MOPC 315, 741F8, 520C9, McPC 603, D1.3, murine phOx, human phOx, RFL3.8 sTCR, 1A6, Se155-4, 18-2-3, 4-4-20, 7A4-1, B6.2, CC49, 3C2, 2c, MA-15C5/ $K_{12}G_0$, Ox, etc. (*see*, Huston, J.S. *et al.*, *Proc. Natl. Acad. Sci. USA* 85:5879-5883 (1988); Huston, J.S. *et al.*, *SIM News* 38(4) (Supp.):11 (1988); McCartney, J. *et al.*, *ICSU Short Reports* 10:114 (1990); McCartney, J.E. *et al.*, unpublished results (1990); Nedelman, M.A. *et al.*, *J. Nuclear Med.* 32 (Supp.):1005 (1991); Huston, J.S. *et al.*, In: *Molecular Design and Modeling: Concepts and Applications, Part B*, edited by J.J. Langone, *Methods in Enzymology* 203:46-88 (1991); Huston, J.S. *et al.*, In: *Advances in the Applications of Monoclonal Antibodies in Clinical Oncology*, Epenetos, A.A. (Ed.), London, Chapman & Hall (1993); Bird, R.E. *et al.*, *Science* 242:423-426 (1988); Bedzyk, W.D. *et al.*, *J. Biol. Chem.* 265:18615-18620 (1990); Colcher,

- D. *et al.*, *J. Nat. Cancer Inst.* 82:1191-1197 (1990); Gibbs, R.A. *et al.*, *Proc. Natl. Acad. Sci. USA* 88:4001-4004 (1991); Milenic, D.E. *et al.*, *Cancer Research* 51:6363-6371 (1991); Pantoliano, M.W. *et al.*, *Biochemistry* 30:10117-10125 (1991); Chaudhary, V.K. *et al.*, *Nature* 339:394-397 (1989); Chaudhary, V.K. *et al.*, *Proc. Natl. Acad. Sci. USA* 87:1066-1070 (1990); Batra, J.K. *et al.*, *Biochem. Biophys. Res. Comm.* 171:1-6 (1990); Batra, J.K. *et al.*, *J. Biol. Chem.* 265:15198-15202 (1990); Chaudhary, V.K. *et al.*, *Proc. Natl. Acad. Sci. USA* 87:9491-9494 (1990); Batra, J.K. *et al.*, *Mol. Cell. Biol.* 11:2200-2205 (1991); Brinkmann, U. *et al.*, *Proc. Natl. Acad. Sci. USA* 88:8616-8620 (1991); Seetharam, S. *et al.*, *J. Biol. Chem.* 266:17376-17381 (1991); Brinkmann, U. *et al.*, *Proc. Natl. Acad. Sci. USA* 89:3075-3079 (1992); Glockshuber, R. *et al.*, *Biochemistry* 29:1362-1367 (1990); Skerra, A. *et al.*, *Bio/Technol.* 9:273-278 (1991); Pack, P. *et al.*, *Biochemistry* 31:1579-1534 (1992); Clackson, T. *et al.*, *Nature* 352:624-628 (1991); Marks, J.D. *et al.*, *J. Mol. Biol.* 222:581-597 (1991); Iverson, B.L. *et al.*, *Science* 249:659-662 (1990); Roberts, V.A. *et al.*, *Proc. Natl. Acad. Sci. USA* 87:6654-6658 (1990); Condra, J.H. *et al.*, *J. Biol. Chem.* 265:2292-2295 (1990); Laroche, Y. *et al.*, *J. Biol. Chem.* 266:16343-16349 (1991); Holvoet, P. *et al.*, *J. Biol. Chem.* 266:19717-19724 (1991); Anand, N.N. *et al.*, *J. Biol. Chem.* 266:21874-21879 (1991); Fuchs, P. *et al.*, *Bio/Technol.* 9:1369-1372 (1991); Breitling, F. *et al.*, *Gene* 104:104-153 (1991); Seehaus, T. *et al.*, *Gene* 114:235-237 (1992); Takkinen, K. *et al.*, *Protein Engng.* 4:837-841 (1991); Dreher, M.L. *et al.*, *J. Immunol. Methods* 139:197-205 (1991); Mottez, E. *et al.*, *Eur. J. Immunol.* 21:467-471 (1991); Traunecker, A. *et al.*, *Proc. Natl. Acad. Sci. USA* 88:8646-8650 (1991); Traunecker, A. *et al.*, *EMBO J.* 10:3655-3659 (1991); Hoo, W.F.S. *et al.*, *Proc. Natl. Acad. Sci. USA* 89:4759-4763 (1993)).

Linkers of the invention used to construct sFv (SCA) polypeptides are designed to span the C-terminus of V_L (or neighboring site thereof) and the N-terminus of V_H (or neighboring site thereof) or between the C-terminus of V_H and the N-terminus of V_L. The preferred length of the peptide linker should be from

2 to about 50 amino acids. In each particular case, the preferred length will depend upon the nature of the polypeptides to be linked and the desired activity of the linked fusion polypeptide resulting from the linkage. Generally, the linker should be long enough to allow the resulting linked fusion polypeptide to properly
5 fold into a conformation providing the desired biological activity. Where conformational information is available, as is the case with sFv (SCA) polypeptides discussed below, the appropriate linker length may be estimated by consideration of the 3-dimensional conformation of the substituent polypeptides and the desired conformation of the resulting linked fusion polypeptide. Where
10 such information is not available, the appropriate linker length may be empirically determined by testing a series of linked fusion polypeptides with linkers of varying lengths for the desired biological activity. Such linkers are described in detail in WO 94/12520, incorporated herein by reference.

Preferred linkers used to construct sFv (SCA) polypeptides have between
15 10 and 30 amino acid residues. The linkers are designed to be flexible, and it is recommended that an underlying sequence of alternating Gly and Ser residues be used. To enhance the solubility of the linker and its associated single chain Fv protein, three charged residues may be included, two positively charged lysine residues (K) and one negatively charged glutamic acid residue (E). Preferably,
20 one of the lysine residues is placed close to the N-terminus of V_H , to replace the positive charge lost when forming the peptide bond of the linker and the V_H . Such linkers are described in detail in U.S. Patent Application S. N. 08/224,591, filed April 7, 1994, incorporated herein by reference. *See also*, Whitlow, M., *et al.*, *Protein Engng.* 7:1017-1026 (1994).

25 For multivalent sFvs (SCA), the association of two or more sFvs (SCA) is required for their formation. Although, multivalent sFvs (SCA) can be produced from sFvs (SCA) with linkers as long as 25 residues, they tend to be unstable. Holliger, P., *et al.*, *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993), have recently demonstrated that linkers 0 to 15 residues in length facilitate the
30 formation of divalent Fvs. *See*, Whitlow, M., *et al.*, *Protein Engng.* 7:1017-1026

(1994); Hoogenboom, H.R., *Nature Biotech.* 15:125-126 (1997); and WO 93/11161.

An object of the present invention is to produce a single-chain antigen-binding polypeptide-polyalkylene oxide conjugate which retains antigen binding
5 affinity within a range of about two-fold to about ten-fold of the antigen binding affinity of the native single-chain antigen-binding polypeptide.

Another object of the present invention is to produce an sFv (SCA) having one or more Cys residues such that the Cys residue is capable of being conjugated with PEG and the PEGylated polypeptide is capable of binding an antigen (i.e., the
10 PEGylated polypeptide's ability to bind an antigen is not disrupted). A further object of the present invention is to produce an sFv (SCA) having three or more consecutive Lys residues such that the Lys residues are capable of being conjugated with PEG and the PEGylated polypeptide is capable of binding an antigen (i.e., the PEGylated polypeptide's ability to bind an antigen is not
15 disrupted). These novel sFv (SCA) proteins may be conjugated to activated polyethylene glycol (PEG) such that the PEG modification occurs only (or preferentially) at the specifically engineered sites. The activated PEG molecules would be thiol-reactive or amine-reactive polymers such as are well known in the art. The designed changes correspond to amino acid residues on the sFv (SCA)
20 surface which are well separated spatially from the antigen-binding site as deduced from known three-dimensional models of the antibody Fv domain.

A further object of the invention is to produce monovalent and multivalent sFvs (SCA) having one or more Cys PEG conjugation sequence(s). A further object of the invention is to produce monovalent and multivalent sFvs (SCA)
25 having three or more consecutive Lys (i.e., oligo-Lys) PEG conjugation sequence(s). For multivalent sFv (SCA), the association of two or more sFvs (SCAs) is required for their formation. For example, multivalent sFvs (SCAs) may be generated by chemically crosslinking two sFvs (SCAs) with C-terminal cysteine residues (Cumber *et al.*, *J. Immunol.* 149:120-126 (1992)) and by linking
30 two sFvs (SCAs) with a third polypeptide linker to form a dimeric Fv

(SCA)(George *et al.*, *J. Cell. Biochem.* 15E:127 (1991)). Details for producing multivalent sFvs (SCAs) by aggregation are described in Whitlow, M., *et al.*, *Protein Engng.* 7:1017-1026 (1994). Multivalent antigen-binding fusion proteins of the invention can be made by any process, but preferably according to the process for making multivalent antigen-binding proteins set forth in WO 93/11161, incorporated herein by reference.

Identification and Synthesis of Site Specific PEGylation Sequences

In the present invention, Cys PEGylation sites may occur in the V_L and V_H regions, adjacent to the C-terminus of the polypeptide (V_L, V_H or neighboring site thereof), the N-terminus of the polypeptide (V_L, V_H or neighboring site thereof), the linker region between the first and second polypeptide regions, or occur in a combination of these regions. In the present invention, oligo-Lys PEGylation sites may occur in the polypeptide linker or in the C-terminus or adjacent to the C-terminus of the polypeptide. The design of the PEG conjugation sites on a protein involves examining the structural information known about the protein and the residues in the proteins involved in antigen binding. The PEG conjugation sites are chosen to be as far from these residues as possible so as to prevent disruption of the antigen-binding site.

The Cys or the oligo-Lys PEGylation site may occur in (1) the native C-terminus of V_L (or V_H), (2) the C-terminus of V_L (or V_H) wherein the C-terminus has a deletion of one or plurality of amino acid residue(s), such that the remaining N-terminus amino acid residues of the peptide are sufficient for the PEGylated polypeptide to be capable of binding an antigen or (3) the C-terminus of V_L (or V_H) wherein the C-terminus has an addition of one or plurality of amino acid residue(s), such that the remaining N-terminus amino acid residues of the peptide are sufficient for the PEGylated polypeptide to be capable of binding an antigen. By "native" is intended the naturally occurring C-terminus of the immunoglobulin (first or second polypeptide). By "C-terminus" it is well understood in the art as

intending the C-terminal amino acid residue or the C-terminal region of the polypeptide, which could include up to all of the amino acid residues of the polypeptide excluding the first N-terminal amino acid residue of the polypeptide. However, in the present invention, "C-terminus" is intended as the C-terminal amino acid residue of the above mentioned three types of C-terminus (1, 2, or 3),
5 unless otherwise indicated or intended.

PEGylation sites were identified and engineered at residues within loop sites in regions of the sFv (SCA) that are diametrically opposed to the antigen binding site. The five loop regions and C-terminal extension chosen as preferred
10 sites of glycosylation are among the most distant regions spatially removed from the binding site.

The six furthest portions of an sFv (SCA) from the antigen binding site are as follows:

- 1) The loop made up of residues 11 to 15 in the light chain;
- 15 2) The loop made up of residues 77 to 79 in the light chain;
- 3) The N-terminus of the linker;
- 4) The loop made up of residues 11 to 15 in the heavy chain;
- 5) The loop made up of residues 82B, 82C and 83 in the heavy chain;
and
- 20 6) The C-terminus of the sFv (or SCA).

The residues are identified as according to Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th ed., U.S. Dept. Health and Human Services, Bethesda, MD (1991). These possible PEGylation sites were determined by examining the 4-4-20 mouse Fab structure (*see*, Whitlow, M. *et al.*, *Protein Engng.* 8:749-761 (1995), incorporated herein by reference).
25

After identifying the loops furthest from the antigen binding site, the nucleic and amino acid sequences of each loop are examined for possible Cys PEGylation sites that may be engineered into the loop region. The engineered placement of the Cys residue anywhere in these six identified regions can generate
30 a preferred site for sFv (SCA) PEGylation. The engineered placement of the

oligo-Lys residues in the linker, the C-terminus of the sFv (SCA) and/or adjacent to the C-terminus of the sFv (SCA) can generate a preferred site for sFv PEGylation.

The design approach described above has been used for the CC49/218 SCA. Figure 2 shows the following resulting designs: designed PEGylation site no. 1 in the light chain of the CC49/218 SCA; designed PEGylation site no. 2 in the N-terminal end of the linker in CC49/218 SCA; designed PEGylation site no. 3 in the heavy chain of the CC49/218 SCA; designed PEGylation site no. 4 at the C-terminus of the CC49/218 SCA. Figure 3 shows the following resulting designs: designed oligo-Lys "hot spot" PEGylation sites at the C-terminus of the CC49/218 SCA. Any combination of these sites could be used.

The particular nucleotide sequence which is used to introduce a Cys or oligo-Lys PEGylation site into the various positions will depend upon the naturally-occurring nucleotide sequence. The most preferred sites are those in which it takes a minimum number of changes to generate the PEGylation site. Of course, based on the redundancy of the genetic code, a particular amino acid may be encoded by multiple nucleotide sequences.

Site-directed mutagenesis is used to change the native protein sequence to one that incorporates the Cys residue or oligo-Lys residues for PEGylation. The mutant protein gene is placed in an expression system, such as bacterial cells, yeast or other fungal cells, insect cells or mammalian cells. The mutant protein can be purified by standard purification methods.

Oligonucleotide-directed mutagenesis methods for generating the Cys or oligo-Lys PEGylation sites and related techniques for mutagenesis of cloned DNA are well known in the art. See, Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989); Ausubel *et al.* (eds.), CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley and Sons (1987), both incorporated herein by reference. A preferred oligonucleotide-directed mutagenesis method for the present invention is according to Ho *et al.*, *Gene* 77:51-59 (1989), incorporated herein by reference.

Hosts and Vectors

After mutating the nucleotide sequence of the sFv (SCA), the mutated DNA can be inserted into a cloning vector for further analysis, such as for confirmation of the DNA sequence. To express the polypeptide encoded by the mutated DNA sequence, the DNA sequence is operably linked to regulatory sequences controlling transcriptional expression and introduced into either a prokaryotic or eukaryotic host cell.

Although sFvs (SCAs) are typically produced by prokaryotic host cells, eukaryotic host cells are the preferred host cells. Preferred host cells include yeast or other fungal cells, insect cells or mammalian cells. Standard protein purification methods may be used to purify these mutant proteins. Only minor modification to the native protein's purification scheme may be required.

Also provided by the invention are DNA molecules such as purified genetic sequences or plasmids or vectors encoding the sFv (SCA) of the invention that have engineered Cys residues and/or oligo-Lys residues capable of PEG conjugation. The DNA sequence for the PEGylated sFv (SCA) polypeptide can be chosen so as to optimize production in organisms such as prokaryotes, yeast or other fungal cells, insect cells or mammalian cells.

The DNA molecule encoding an sFv (SCA) having Cys residues and/or oligo-Lys residues for PEG conjugation can be operably linked into an expression vector and introduced into a host cell to enable the expression of the engineered sFv (SCA) protein by that cell. A DNA sequence encoding an sFv (SCA) having Cys and/or oligo-Lys PEGylation sites may be recombined with vector DNA in accordance with conventional techniques. Recombinant hosts as well as methods of using them to produce single chain proteins of the invention are also provided herein.

The expression of such sFv (SCA) proteins of the invention can be accomplished in procaryotic cells. Preferred prokaryotic hosts include, but are not

limited to, bacteria such as *Neisseria*, *Mycobacteria*, *Streptococci*, *Chlamydia* and *E. coli*.

Eukaryotic hosts for cloning and expression of such sFv (SCA) proteins of the invention include insect cells, yeast, fungi, and mammalian cells (such as, 5 for example, human or primate cells) either *in vivo*, or in tissue culture. A preferred host for the invention is *Pichia pastoris*.

The appropriate DNA molecules, hosts, methods of production, isolation and purification of monovalent, multivalent and fusion forms of proteins, especially sFv (SCA) polypeptides, are thoroughly described in the prior art, such 10 as, e.g., U.S. Patent No. 4,946,778, which is fully incorporated herein by reference.

The sFv (SCA) encoding sequence having Cys residues and/or oligo-Lys residues for PEG conjugation and an operably linked promoter may be introduced into a recipient prokaryotic or eukaryotic cell either as a non-replicating DNA (or 15 RNA) molecule, which may either be a linear molecule or, more preferably, a closed covalent circular molecule. Since such molecules are incapable of autonomous replication, the expression of the desired sFv (SCA) protein may occur through the transient expression of the introduced sequence. Alternatively, permanent expression may occur through the integration of the introduced sFv 20 (SCA) sequence into the host chromosome.

In one embodiment, the sFv (SCA) sequence can be integrated into the host cell chromosome. Cells which have stably integrated the introduced DNA into their chromosomes can be selected by also introducing one or more markers which allow for selection of host cells which contain the sFv (SCA) sequence and 25 marker. The marker may complement an auxotrophy in the host (such as *his4*, *leu2*, or *ura3*, which are common yeast auxotrophic markers), biocide resistance, e.g., antibiotics, or resistance to heavy metals, such as copper, or the like. The selectable marker gene can either be directly linked to the sFv (SCA) DNA sequence to be expressed, or introduced into the same cell by co-transfection.

In another embodiment, the introduced sequence will be incorporated into a plasmid vector capable of autonomous replication in the recipient host cell. Any of a wide variety of vectors may be employed for this purpose. Factors of importance in selecting a particular plasmid or viral vector include: the ease with which recipient cells that contain the vector may be recognized and selected from those recipient cells which do not contain the vector; the number of copies of the vector which are desired in a particular host; and whether it is desirable to be able to "shuttle" the vector between host cells of different species.

Any of a series of yeast vector systems can be utilized. Examples of such expression vectors include the yeast 2-micron circle, the expression plasmids YEP13, YCP and YRP, etc., or their derivatives. Such plasmids are well known in the art (Botstein *et al.*, *Miami Wntr. Symp.* 19:265-274 (1982); Broach, J.R., In: *The Molecular Biology of the Yeast Saccharomyces: Life Cycle and Inheritance*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, p. 445-470 (1981); Broach, J.R., *Cell* 28:203-204 (1982)).

For a mammalian host, several possible vector systems are available for expression. One class of vectors utilize DNA elements which provide autonomously replicating extra-chromosomal plasmids, derived from animal viruses such as bovine papilloma virus, polyoma virus, adenovirus, or SV40 virus. A second class of vectors relies upon the integration of the desired gene sequences into the host chromosome. Cells which have stably integrated the introduced DNA into their chromosomes may be selected by also introducing one or more markers which allow selection of host cells which contain the expression vector. The marker may provide for prototrophy to an auxotrophic host, biocide resistance, e.g., antibiotics, or resistance to heavy metals, such as copper or the like. The selectable marker gene can either be directly linked to the DNA sequences to be expressed, or introduced into the same cell by co-transformation. Additional elements may also be needed for optimal synthesis of mRNA. These elements may include splice signals, as well as transcription promoters, enhancers, and termination signals. The cDNA expression vectors incorporating such


elements include those described by Okayama, H., *Mol. Cell. Biol.* 3:280 (1983), and others.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript
 5 vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Preferred vectors for expression in *Pichia* are pHIL-S1
 10 (Invitrogen Corp.) and pPIC9 (Invitrogen Corp.). Other suitable vectors will be readily apparent to the skilled artisan.

Once the vector or DNA sequence containing the constructs has been prepared for expression, the DNA constructs may be introduced or transformed into an appropriate host. Various techniques may be employed, such as
 15 transformation, transfection, protoplast fusion, calcium phosphate precipitation, electroporation, or other conventional techniques. After the cells have been transformed with the recombinant DNA (or RNA) molecule, the cells are grown in media and screened for appropriate activities. Expression of the sequence results in the production of the mutant sFv (SCA) for PEG conjugation of the
 20 present invention.

Straight chain Polymers

The straight chain polyalkylene glycols employed in the practice of the present invention are of the structural formula



25 wherein R is selected from the group consisting of hydrogen, lower alkyl, and mixtures thereof, R¹ is selected from the group consisting of hydrogen and lower

alkyl, and n is a positive integer. By "lower alkyl" is meant an alkyl group having from one to four carbon atoms, i.e., methyl, ethyl, propyl, butyl, and isomers of the foregoing. R is preferably selected from the group consisting of hydrogen, methyl, and mixtures thereof, R¹ is preferably selected from the group consisting of hydrogen and methyl, and n is preferably a positive integer of 500 or less. R is most preferably hydrogen, R¹ is most preferably methyl, and n is most preferably an integer of 7 to 150. It will be readily apparent to those skilled in the art that the preferred poly(alkylene glycols) employed in the practice of the present invention are poly(ethylene glycol), poly(propylene glycol), mixtures thereof, and copolymers of poly(ethylene glycol) and poly(propylene glycol), wherein one of the terminal hydroxyl groups of the polymer may be substituted with a lower alkyl group. A preferred polyalkylene glycol for use in the present invention is poly(ethylene glycol)-hydrazine. The most preferred polyalkylene glycol for use in the present invention is methoxy poly(ethylene glycol).

Hereinafter, for convenience, the polyalkylene glycol employed in the practice of the present invention will be designated PAG, which term is intended to include both compounds wherein R¹ is hydrogen and compounds wherein R¹ is alkyl. PEG refers to poly(ethylene glycol) and mPEG refers to methoxy poly(ethylene glycol).

The PAG does not have to be of a particular molecular weight, but it is preferred that the molecular weight be between about 500 and about 40,000; more preferably, between about 2,000 and about 20,000. The choice of molecular weight of PAG is made based on the nature of the particular polypeptide employed, for example, the number of amino or other groups available on the polypeptide for modification. Molecular weights of about 10,000 and about 20,000 are most preferred.


It is well known in the art that PAGs that contain two terminal hydroxyl groups per moiety are capable of crosslinking other polymers, e.g. proteins. Where, as is often the case, crosslinking would be deemed undesirable, such crosslinking can be minimized or prevented by means known in the art. For

example, Davis *et al.* in U.S. Patent No. 4,179,337 have pointed out that a preferred means for preventing crosslinking is to preblock one end of the PAG, such as is done in the commercially available methoxy poly(ethylene glycol).

The PAGs employed in the practice of the present invention are preferably coupled to polypeptides by means of suitable coupling agents. A useful review of a number of coupling agents that can be employed in the practice of the present invention appears in Dreborg *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems* 6(4):315-365 (1990), see, especially, pp. 317-320.

Probably the best known coupling agent for this purpose is cyanuric chloride. Its use has been described in numerous references, see, for example, Abuchowski *et al.*, *J. Biol. Chem.* 252(11):3578-3581 (June 10, 1977).

Zalipsky *et al.*, *Eur. Pol. J.* 19(12):1177-1183 (1983), among others, have described the reaction of methoxy poly(ethylene glycol) with succinic anhydride:




It is also known to alkylate mPEG with ethylbromoacetate in the presence of a base such as K-tertiary butoxide in tertiary butanol, Na-naphthalene in tetrahydrofuran, or butyl lithium in benzene:




The terminal hydroxyl groups of PEG can be transformed into amine, carboxyl, or hexamethyl isocyanate groups. See, for example, Zalipsky *et al.*, 1983, *supra*. A mixed anhydride derivative of carboxylated mPEG can be prepared in the presence of triethylamine and then reacted with proteins:


-34-




Carboxylated mPEG can also be reacted with hydroxysuccinimide in the presence of dicyclohexylcarbodiimide and dimethyl formamide for reaction with protein:





- 5 King and Weiner (*Int. J. Peptide Protein Res.* 16:147 (1980)) describe the dithiocarbonate of mPEG:



Beauchamp *et al.*, *Analytical Biochem.* 131:25-33 (1983) describe the activation of PEG with 1,1'-carbonyldiimidazole. Reaction of this derivative with a peptide yields a carbamate linkage:




- 5 Veronese *et al.*, *Appl. Biochem. & Biotechnol.* 11:141-152 (1985) describe the activation of methoxy poly(ethylene glycol) with phenylchloroformates, e.g., 2,4,5-trichlorophenylchloroformate or *p*-nitrophenylchloroformate. These derivatives are linked to peptides by urethane linkages:




- 10 Ueno *et al.* in European Patent Application 87103259.5 form mPEG imidoesters from the corresponding nitriles by reaction with dry hydrogen chloride in the presence of a dehydrated lower alcohol:

-36-




Abuchowski *et al.*, *Cancer Biochem. Biophys.* 7:175-186 (1984) have described forming mPEG succinate as described above and then forming methoxy polyethylene glycolyl succinimidyl succinate ("SS-PEG") by reaction with hydroxysuccinimide in the presence of dicyclohexylcarbodiimide:




-37-

Sano *et al.*, European Patent Application No. 89107960.0 disclose the phenyl glyoxal derivative of methoxy poly(ethylene glycol), which is capable of modifying the guanidino groups in peptides:




Zalipsky, in U.S. Patent No. 5,122,614, describes the activation of PEG
 5 by conversion into its N-succinimide carbonate derivative ("SC-PEG"):



methoxypoly(ethylene glycol) - succinyl carbonate


SC - PEG

Zalipsky *et al.*, *J. Macromol. Sci. Chem. A21*:839, disclose the amino acid ester derivative of methoxy poly(ethylene glycol):



-38-

Davis *et al.*, U.S. Patent No. 4,179,337, disclose a hydrazide derivative of methoxy poly(ethylene glycol), which is capable of modifying aldehydes and ketones and other functional groups:




5 It is further disclosed that the bifunctional derivative of PEG, i.e., polyethylene glycol-bis-succinidyl carbonate ("BSC-PEG") can be prepared by similar means. The SC-PEG and BSC-PEG compounds are then reacted with amine groups in a protein and attached thereto via urethane (carbamate) linkages.

10 It will be readily apparent to those skilled in the art that other activated PAGs can also be employed in the practice of the present invention. The preferred activated PAG for use in the practice of the present invention is selected from the group consisting of SS-PEG and SC-PEG. The use of SC-PEG is most preferred.

Branched Polymers

15 The invention further provides for the use of branched, substantially non-antigenic polymers for polyalkylene oxide conjugation of the sFv (SCA) proteins corresponding to the formula:



wherein (R) includes a water-soluble non-antigenic polymer;

(n) = 2 or 3;

20 (L) is an aliphatic linking moiety covalently linked to each (R); and (A) represents an activated functional group capable of undergoing nucleophilic substitution. For example, (A) can be a group which is capable of bonding with biologically active nucleophiles or moieties capable of doing the same.

25 In particularly preferred aspects of the invention (R) includes a poly(alkylene oxide) PAO such as poly(ethylene glycol) PEG or mPEG. It is

preferred that each chain have a molecular weight of between about 200 and about 12,000 daltons and preferably between about 1,000 and about 10,000 daltons. Molecular weights of about 5,000 daltons are most preferred.

As shown in Formula II, 2 or 3 polymer chains, designated (R) herein, are
5 joined to the alphatic linking moiety (L). Suitable aliphatics included substituted alkyl diamines and triamines, lysine esters and malonic ester derivatives. The linking moieties are perferably non-planar, so that the polymer chains are not rigidly fixed. The linking moiety (L) is also a means for attaching the mulitple polymer chains or "branches" to (A), the moeity through which the polymer
10 attaches to the sFv (SCA) protein.

(L) preferably includes a multiply-functionalized alykyl group containing up to 18, and more preferably between 1-10 carbon atoms. A heteroatom such as nitrogen, oxygen or sulfur may be included within the alkyl chain. The alkyl chain may also be branched at a carbon or nitrogen atom. In another aspect of the
15 invention, (L) is a single nitrogen atom.

(L) and (R) are preferably joined by a reaction between nucleophilic functional groups on both (R) and (L). Each (R) is suitably functionalized to undergo nucleophilic substitution and bond with (L). Such functionalization of polymers is readily apparent to those of ordinary skill in the art.

20 A wide variety of linkages are contemplated between (R) and (L). Urethane (carbamate) linkages are preferred. The bond can be formed, for example, by reacting an amino group such as 1,3-diamino-2-propanol with methoxypolyethylene glycol succinimidyl carbonate as described in U.S. Patent No. 5,122,614. Amide linkages, which can be formed by reacting an amino-
25 terminated non-antigenic polymer suchas methoxypolyethylene glycol-amine (mPEG amine) with an acyl chloride functional group. Examples of other such linkages include ether, amine, urea, and thio and thiol analogs thereof, as well as the thio and thiol analogs of the urethane and amide linkages discussed supra.

The moiety (A) of Formula II represents groups that "activate" the branched polymers of the present invention for conjugation with biologically active materials. (A) can be a moiety selected from:

1. Functional groups capable of reacting with an amino group such as:
 - 5 a) carbonates such as the p-nitrophenyl or succinimidyl;
 - b) carbonyl imidazole;
 - c) azlactones;
 - d) cyclic imide thiones; or
 - e) isocyanates or isothiocyanates.
- 10 2. Functional groups capable of reacting with carboxylic acid groups and reactive with carbonyl groups such as:
 - a) primary amines; or
 - b) hydrazine and hydrazide functional groups such as the acyl hydrazides, carbazates, semicarbamates, thiocarbazates, etc.
- 15 3. Functional groups capable of reacting with mercapto or sulfhydryl groups such as phenyl glyoxals; see, for example, U.S. Patent No. 5,093,531.
4. Other nucleophiles capable of reacting with an electrophilic center. A non-limiting list includes, for example, hydroxyl, amino, carboxyl, thiol groups, active methylene and the like.

20 The moiety (A) can also include a spacer moiety located proximal to the aliphatic linking moiety (L). The spacer moiety may be a heteroalkyl, alkoxy, alkyl containing up to 18 carbon atoms or even an additional polymer chain. The spacer moieties can be added using standard synthesis techniques.

25 The branched polymers, generally, U-PAO's or U-PEG's, are formed using conventional reaction techniques known to those of ordinary skill in the art.


 These umbrella-like branched polymers of the present invention (U-PAO's or U-PEG's) react with biologically active nucleophiles to form conjugates. The point of polymer attachment depends upon the functional group (A). For example, (A) can be a succinimidyl succinate or carbonate and react with ϵ -amino lysines.

30 The branched polymers can also be activated to link with any primary or

secondary amino group, mercapto group, carboxylic acid group, reactive carbonyl group or the like found on biologically active polypeptides. Other groups are apparent to those of ordinary skill in the art.

One of the main advantages of the use of the branched polymers is that the branching imparts an umbrella-like three dimensional protective covering to the materials they are conjugated with. This contrasts with the string-like structure of the straight chain polymers discussed, supra. An additional advantage of the branched polymers is that they provide the benefits associated with attaching several strands of polymers to a sFv protein but require substantially fewer conjugation sites. The desired properties of PEGylation are realized and the loss of bioactivity is minimized.

One or more of the activated branched polymers can be attached to a biologically active nucleophile, such as an sFv protein, by standard chemical reactions. The conjugate is represented by the formula:



wherein (R) is a water-soluble substantially non-antigenic polymer; $n=2$ or 3 ; (L) is an aliphatic linking moiety; (A^1) represents a linkage between (L) and the nucleophile and (z) is an integer ≥ 1 representing the number of polymers conjugated to the biologically active nucleophile. The upper limit for (z) will be determined by the number of available nucleophilic attachment sites and the degree of polymer attachment sought by the artisan. The degree of conjugation can be modified by varying the reaction stoichiometry using well-known techniques. More than one polymer conjugated to the nucleophile can be obtained by reacting a stoichiometric excess of the activated polymer with the nucleophile.

Purification of sFv Proteins

A generic protocol that has been developed and used to produce twelve different single chain antigen binding molecules. It involves cell lysis and washing, solubilization in a denaturing solvent, refolding by dilution, and two ion-exchange
5 HPLC chromatography steps. Such isolated sFvs (SCAs) are capable of being PAG conjugated according to the present invention.

The fermentation of the sFv-producing *E. coli* strains are performed at 32°C using a casein digest-glucose-salts medium. At an optical density of 18 to 20 at 600 nm, sFv expression is induced by a 42°C temperature shock for one
10 hour. After the fermentation is cooled to 10°C, the cells are harvested by centrifugation at 7000g for ten minutes. The wet cell paste is then stored frozen at -20°C. Approximately 200 to 300 g of wet cell paste is normally recovered from one 10-liter fermentation.

For protein recovery, the cell paste from three 10-liter fermentations (600-
15 900 g) is thawed overnight at 4°C and gently resuspended at 4°C in 50 mM Tris-HCl, 1.0 mM EDTA, 100 mM KCl, 0.1 mM phenylmethylsulfonyl chloride (PMSF), pH 8.0 (lysis buffer), using 10 liters of lysis buffer for every kilogram of wet cell paste. When thoroughly resuspended, the chilled mixture is passed three times through a Manton-Gaulin cell homogenizer to fully lyse the cells. Because
20 the cell homogenizer raises the temperature of the cell lysate to 25±5°C, the cell lysate is cooled to 5±2°C with a Lauda/Brinkman chilling coil after each pass. Complete lysis is verified by visual inspection under a microscope.

The cell lysate is centrifuged at 24,300g for thirty minutes at 6°C using a Sorvall RC-5B centrifuge. The pellet contains the insoluble sFv and the
25 supernatant is discarded. The pellet is washed by gently scraping it from the centrifuge bottles and resuspending it in 5 liters of lysis buffer/kg of wet cell paste. The resulting 3.0-4.5-liter suspension is again centrifuged at 24,300g for 30 min at 6°C, and the supernatant is discarded. This washing of the cell pellet removes soluble *E. coli* proteins and can be repeated as many as five times. At any time

-43-

during this washing procedure the material can be stored as a frozen pellet at -20°C. A substantial time saving in the washing steps can be accomplished by utilizing a Pellicon tangential flow apparatus equipped with 0.22-µm microporous filters.

5 The washed cell pellet is solubilized at 4°C in freshly prepared 6 M guanidine hydrochloride, 50 mM Tris-HCl, 10 mM CaCl₂, 50 mM KCl, pH 8.0 (denaturing buffer), using 6 ml/g of pellet. If necessary, a few quick pulses from a Heat Systems Ultrasonics tissue homogenizer can be used to complete the solubilization. The resulting suspension is centrifuged at 24,300g for 45 minutes
10 at 6°C and the pellet is discarded. The optical density of the supernatant is determined at 280 nm and if the OD₂₈₀ is above 30, additional denaturing buffer is added to obtain an OD₂₈₀ of approximately 25.

 The supernatant is slowly diluted into cold (4-7°C) refolding buffer (50 mM Tris-HCl, 10 mM CaCl₂, 50 mM KCl, 0.1 mM PMSF, pH 8.0) until a 1:10
15 to 1:100 dilution is reached (final volume 70-120 liters). The refolding buffer should be prepared at least one day prior to use, to allow sufficient time for it to cool to 4°C. The best results will be obtained when the supernatant is slowly added to the refolding buffer over a two hour period, with gentle mixing. The solution is left undisturbed for at least twenty hours and then filtered through a
20 Millipore Pellicon tangential flow apparatus at 4°C with four to six 0.45-µm microporous membranes (HVLP 000 C5). The filtrate is concentrated to 1 to 2 liters using a Pellicon apparatus with four to six 10,000 NMWL cassettes (SK1PA156A4), again at 4°C.

 The concentrated crude sFv sample is buffer exchanged at 4°C into 20
25 mM 2-[N-morpholino]ethanesulfonic acid (Mes), 0.3 mM CaCl₂, pH 6.0, using the Pellicon ultrafiltration apparatus equipped with four to six 10,000 NMWL cassettes. The sample is then chromatographed on a Waters Accell Plus CM ion-exchange (RCM) column (4.7 x 30.0 cm). Prior to loading on the HPLC, the material is filtered through a 0.22-µm filter and the Accell column is equilibrated
30 with Buffer A (40 mM Mes, 1 mM CaCl₂, pH 6.0). Following sample loading, the

Accell column is eluted over a 55-minute period with a linear gradient of Buffer A and Buffer B (40 mM Mes, 100 mM CaCl₂, pH 7.0). (See Table 1).

Table 1				
Accell Cation-Exchange HPLC Gradients				
Time (min) ^a	Flow (ml/min)	Buffers ^b		
		%A	%B	%C
Initial	40.0	100	0	0
55.0	40.0	0	100	0
58.0	40.0	0	100	0
60.0	40.0	0	0	100
62.0	40.0	100	0	0
^a Linear gradients are run between each time point.				
^b Buffer A, 40 mM Mes, 1 mM CaCl ₂ , pH 6.0; Buffer B, 40 mM Mes, 100 mM CaCl ₂ , pH 7.0; Buffer C, 40 mM Mes, 20% ethanol, pH 7.5.				

The Accell Plus CM column has a capacity of about 3 g and thus all the crude sFv sample can normally be loaded in a single run. The fractions are analyzed using 4-20% Novex SDS-PAGE gels and the peak fractions are pooled. Normally, the sFv elutes from the Accell ion-exchange column quite early in the gradient. To enhance resolution for certain sFv proteins, holds in the gradient can be implemented.

The pooled fractions from the Accell HPLC purification are dialyzed against Buffer D (40 mM 3-[N-morpholino]propanesulfonic acid (Mops), 0.5 mM Ca acetate, pH 6.0) until the conductivity is lowered to that of Buffer D. The sample is then loaded on a 21.5 x 150-mm polyaspartic acid PolyCAT A column. If more than 60 mg is loaded on this column, the resolution begins to deteriorate; thus, the pooled fractions from the Accell HPLC purification often must be divided into several PolyCAT A runs. Most sFv proteins have an extinction coefficient of about 2.0 mg ml⁻¹ cm⁻¹ at 280 nm and this can be used to determine protein concentration. The sFv sample is eluted from the PolyCAT A column with

a 50-min linear gradient of Buffer D and Buffer E (40 mM Mops, 10 mM Ca acetate, pH 8.0). See Table 2.

Table 2				
PolyCAT A Cation-Exchange HPLC Gradients				
Time (min) ^a	Flow (ml/min)	Buffers ^b		
		%D	%E	%F
Initial	15.0	100	0	0
50.0	15.0	0	100	0
55.0	15.0	0	100	0
60.0	15.0	0	0	100
63.0	15.0	0	0	100
64.0	15.0	100	0	0
67.0	15.0	100	0	0
^a Linear gradients are run between each time point.				
^b Buffer D, 40 mM Mops, 0.5 mM Ca acetate, pH 6.0; Buffer E, 40 mM Mops, 10 mM Ca acetate, pH 8.0; Buffer F, 40 mM Mops, 100 mM CaCl ₂ , pH 7.5.				

The sFv proteins will often elute between 20 and 26 min when this gradient is used. This corresponds to an eluting solvent composition of approximately 70% Buffer D and 30% Buffer E.

This purification procedure yields sFv proteins that are more than 95% pure as examined by SDS-PAGE and Scatchard analysis. Modifications of the above procedure may be dictated by the isoelectric point of the particular sFv being purified, which is often between 8.0 and 9.3.

The polyalkylene glycols (PAGs) employed in the practice of the present invention, which, as indicated above, are preferably activated by reaction with a coupler, can be reacted with any of several groups that may be present attached to the chain of the single chain antigen binding molecules, e.g. terminal carboxyl groups, thiol groups, phenolic hydroxyl groups, or primary amino groups located at the chain terminus or along the chain. It is preferred to react activated PAGs with primary amine groups, especially those occurring along the peptide chain.

It is most preferred that the activated PAGs be coupled to the ϵ amino groups of lysine residues as well as cysteine residues in the polypeptide.

The reaction between the PAG and the single chain polypeptide is normally carried out in solution, preferably an aqueous buffer solution providing a pH in the
 5 range of from about 6 to about 10, preferably from about 7 to about 9, most preferably from about 7 to about 8. As examples of buffer solutions that will provide pH's in these ranges at 25° C may be listed:

- 50 ml of 0.1 molar potassium dihydrogen phosphate + 5.6 to 46.1
 ml 0.1 molar NaOH diluted to 100 ml
- 10 50 ml of 0.025 molar borate + 2.0 to 20.5 ml 0.1 molar HCl
 diluted to 100 ml
- 50 ml of 0.025 molar borate + 0.9 to 18.3 ml 0.1 molar NaOH
 diluted to 100 ml
- 50 ml of 0.05 molar sodium bicarbonate + 5.0 to 10.7 ml 0.1
 15 molar NaOH diluted to 100 ml

The precise adjustment of the quantity of acid or base to be used to provide a particular desired pH will be readily determinable by those skilled in the art.

If, in a given instance, the use of a biological buffer should be required, one of the following may be employed:

- 20 3-(N-Morpholino)propanesulfonic acid (MOPS)
- 3-(N-Morpholino)-2-hydroxypropanesulfonic acid (MOPSO)
- Piperazine-N,N'-bis(2-hydroxypropanesulfonic acid) (POPSO)

The reaction between the PAG and the single chain polypeptide will normally be run under conditions that will not give rise to denaturation, e.g. mild
 25 temperatures and no more agitation than necessary. The reaction will preferably be run at a temperature in the range of from about 4° C to about 25° C. More

preferably, the reaction will be run at room temperature, i.e. from about 20° C to about 25° C.

It will be readily understood by those skilled in the art that the amount of PAG employed relative to the amount of single chain polypeptide will be dependent upon the desired nature of the reaction product. Where, for example, it is desired to react a PAG with each lysine residue along the polypeptide chain, an amount of PAG at least equimolar to the lysine concentration will be required. It will be advantageous to employ an excess of PAG, where possible, in order to increase the reaction rate and the likelihood of a complete reaction. Clearly, if fewer than all of the possible reaction sites along the polypeptide chain are to be derivatized, correspondingly less PAG will be used. In general, however, where molar excesses of PAG's are used, it has been determined that molar excesses on the order of 2 - 100 of the PAG can be used; molar excesses of 2 - 10 are preferred.

The time required for the reaction will depend upon a number of factors, such as reaction temperature, concentration of reactants, and whether full or partial reaction is desired. The course of the reaction can be monitored by conventional means, such as the analysis of periodic samples by size exclusion chromatography or gel electrophoresis. The reaction can conveniently be terminated when desired by the addition of a compound having a primary amine group, e.g. glycine, to scavenge the excess PAG. A reaction time of about 15 - 120 minutes will typically be required to fully react the PAG with the primary amine groups of the lysine residues of the single chain polypeptide at room temperature. The skilled practitioner will understand that the time for conjugation, as well as the amount and type of PAG, must not be such as to inactivate the polypeptide being employed.

Purification of the PAG/single chain polypeptide reaction product can be effected by means commonly employed by those skilled in the art, such as, for example, size exclusion chromatography, ion-exchange chromatography, ultrafiltration, dialysis, and the like. Solutions of the reaction product can, if

desired, be concentrated with a rotary evaporator and can be obtained in the dry state by lyophilization.

Depending upon the particular single chain antigen binding molecule chosen and the extent to which it is reacted with the PAG, the resulting adduct is
5 expected to be useful both diagnostically and therapeutically, exhibiting, as compared to the unreacted single chain polypeptide, decreased immunogenicity, increased circulating life, and increased stability while maintaining an acceptable level of activity.

The single chain antigen binding polypeptide can be reacted with the
10 activated branched polyethylene glycol polymers discussed above in an aqueous reaction medium which can be buffered, depending on the pH requirements of the nucleophile. The optimum pH for the reaction is generally between about 6.5 and about 8.0 and preferably about 7.4 for polypeptides. The optimum reaction conditions for the sFv stability, reaction efficiency, etc., is within the level of
15 ordinary skill in the art. The preferred temperature range is between 4°C and 37°C. The reaction temperature cannot exceed the temperature at which the nucleophile may denature or decompose. It is preferred that the nucleophile be reacted with an excess of the activated branched polymer. Following the reaction, the conjugate is recovered and purified, for example, by diafiltration, column
20 chromatography, combinations thereof, or the like.

Conjugates

Upon production of the polyalkylene oxide conjugated sFv (SCA) of the present invention, the polyalkylene oxide conjugated sFv may further be modified by conjugating a diagnostic or therapeutic agent to the polyalkylene oxide
25 conjugated sFv. The general method of preparing an antibody conjugate according to the invention is described in Shih, L.B., *et al.*, *Cancer Res.* 51:4192 (1991); Shih, L.B., and D.M. Goldenberg, *Cancer Immunol. Immunother.* 31:197 (1990); Shih, L.B., *et al.*, *Intl. J. Cancer* 46:1101 (1990); Shih, L.B., *et al.*, *Intl.*

J. Cancer 41:832 (1988), all incorporated herein by reference. The indirect method involves reacting an antibody (or sFv), whose polyalkylene oxide has a functional group, with a carrier polymer loaded with one or plurality of bioactive molecules, such as, peptides, lipids, nucleic acids (i.e., phosphate-lysine
5 complexes), drug, toxin, chelator, boron addend or detectable label molecule(s).

Alternatively, the polyalkylene oxide conjugated sFv may be directly conjugated with a diagnostic or therapeutic agent. The general procedure is analogous to the indirect method of conjugation except that a diagnostic or therapeutic agent is directly attached to an oxidized sFv component. *See Hansen*
10 *et al.*, U.S. Patent No. 5,443,953, incorporated herein by reference.

The polyalkylene oxide conjugated sFv can be attached to a derivative of the particular drug, toxin, chelator, boron addend or label to be loaded, in an activated form, preferably a carboxyl-activated derivative, prepared by conventional means, e.g., using dicyclohexylcarbodiimide (DCC) or a water
15 soluble variant thereof, to form an intermediate adduct.

Many drugs and toxins are known which have a cytotoxic effect on tumor cells or microorganisms that may infect a human and cause a lesion, in addition to the specific illustrations given above. They are to be found in compendia of drugs and toxins, such as the Merck Index and the like. Any such drug can be loaded
20 onto a carrier or directly onto a polyalkylene oxide conjugated sFv by conventional means well known in the art, and illustrated by analogy to those described above.

Chelators for radiometals or magnetic resonance enhancers are also well known in the art. Typical are derivatives of ethylenediaminetetraacetic acid
25 (EDTA) and diethylenetriaminepentaacetic acid (DTPA). These typically have groups on the side chain by which the chelator can be attached to a carrier or directly onto a polyalkylene oxide conjugated sFv. Such groups include, e.g., a benzylisothiocyanate, by which the DTPA or EDTA can be coupled to the reactive group of an sFv.

Labels such as radioisotopes, enzymes, fluorescent compounds, electron transfer agents, and the like can also be linked to carrier or directly onto a polyalkylene oxide conjugated sFv by conventional methods well known to the art. These labels and the sFv conjugates prepared from them can be used for immunoassays and for immunohistology, much as the sFv conjugate prepared by direct attachment of the labels to the sFv. However, the loading of the conjugates according to the present invention with a plurality of labels can increase the sensitivity of assays or histological procedures, where only low extent of binding of the sFv to target antigen is achieved.

Boron addends, e.g., carboranes, when attached to single-chain antigen binding molecules and targeted to lesions, can be activated by thermal neutron irradiation and converted to radioactive atoms which decay by alpha emission to produce highly cytotoxic short-range effects. High loading of boron addends, as well as of magnetic resonance enhancing ions, is of great importance in potentiating their effects. Carboranes can be made with carboxyl functions on pendant side chains, as is well known in the art.

Loading of drugs on the carrier will depend upon the potency of the drug, the efficiency of sFv targeting and the efficacy of the conjugate once it reaches its target. In most cases, it is desirable to load at least 20, preferably 50, and often 100 or more molecules of a drug on a carrier. The ability to partially or completely detoxify a drug as a conjugate according to the invention, while it is in circulation, can reduce systemic side effects of the drug and permit its use when systemic administration of the unconjugated drug would be unacceptable. Administration of more molecules of the drug, but conjugated to the sFv on a carrier, according to the present invention, permits therapy while mitigating systemic toxicity.

Toxins will often be less heavily loaded than drugs, but it will still be advantageous to load at least 5, preferably 10 and in some cases 20 or more molecules of toxin on a carrier and load at least one carrier chain on the sFv for targeted delivery.

Uses

The polyalkylene oxide conjugated sFv (SCA) polypeptide conjugates of the present invention are expected to have much longer circulating half lives and reduced immunogenicity *in vivo*. This may solve a potential limitation relating to very rapid blood clearance of some sFv proteins. It would also reduce or eliminate concerns about repeated administration of a therapeutic sFv which may otherwise provoke an immune response in the patient. The choice of the particular cysteine and/or oligo-lysine mutant combinations may allow one to achieve circulating lives over a considerable range depending on the specific polyalkylene oxide conjugated sFv variant polypeptide. This would allow sFv to be administered for the therapeutic use of choice.

A diagnostic or therapeutic agent is a molecule or atom which is conjugated to an antibody and useful for diagnosis or for therapy. The immunoreactivity of the antibody is retained. Diagnostic or therapeutic agents include drugs, toxins, chelators, boron compounds and detectable labels. *See* "Conjugates" section, *supra*, for further details.

The diagnostic or therapeutic agent may be, but is not limited to, at least one selected from a nucleic acid, a compound, a protein, an element, a lipid, an antibody, a saccharide, an isotope, a carbohydrate, an imaging agent, a lipoprotein, a glycoprotein, an enzyme, a detectable probe, or any combination thereof, which may be detectably labeled as for labeling antibodies, as described herein. Such labels include, but are not limited to, enzymatic labels, radioisotope or radioactive compounds or elements, fluorescent compounds or metals, chemiluminescent compounds and bioluminescent compounds. Alternatively, any other known diagnostic or therapeutic agent can be used in a method of the present invention.

A therapeutic agent used in the present invention may have a therapeutic effect on the target cell, the effect selected from, but not limited to, correcting a defective gene or protein, a drug action, a toxic effect, a growth stimulating effect, a growth inhibiting effect, a metabolic effect, a catabolic affect, an anabolic effect,

an antiviral effect, an antibacterial effect, a hormonal effect, a neurohumoral effect, a cell differentiation stimulatory effect, a cell differentiation inhibitory effect, a neuromodulatory effect, an antineoplastic effect, an anti-tumor effect, an insulin stimulating or inhibiting effect, a bone marrow stimulating effect, a pluripotent stem cell stimulating effect, an immune system stimulating effect, and any other known therapeutic effects that may be provided by a therapeutic agent delivered to a cell via a delivery system according to the present invention.

The sFv conjugate of the present invention may be used for protection, suppression or treatment of infection or disease. By the term "protection" from infection or disease as used herein is intended "prevention," "suppression" or "treatment." "Prevention" involves administration of a glycosylated sFv conjugate prior to the induction of the disease. "Suppression" involves administration of the composition prior to the clinical appearance of the disease.

"Treatment" involves administration of the protective composition after the appearance of the disease. It will be understood that in human and veterinary medicine, it is not always possible to distinguish between "preventing" and "suppressing" since the ultimate inductive event or events may be unknown, latent, or the patient is not ascertained until well after the occurrence of the event or events. Therefore, it is common to use the term "prophylaxis" as distinct from "treatment" to encompass both "preventing" and "suppressing" as defined herein. The term "protection," as used herein, is meant to include "prophylaxis."

Such additional therapeutic agents which can further comprise a therapeutic agent or composition of the present invention may be selected from, but are not limited to, known and new compounds and compositions including antibiotics, steroids, cytotoxic agents, vasoactive drugs, antibodies and other therapeutic modalities. Non-limiting examples of such agents include antibiotics used in the treatment of bacterial shock, such as gentamycin, tobramycin, nafcillin, parenteral cephalosporins, etc; adrenal corticosteroids and analogs thereof, such as methyl prednisolone, mitigate the cellular injury caused by endotoxins; vasoactive drugs, such as alpha receptor blocking agent (e.g., phenoxybenzamine),

beta receptor agonists (e.g., isoproterenol), and dopamine are agents suitable for treating septic shock.

Polyalkylene oxide conjugated sFv of the invention may also be used for diagnosis of disease and to monitor therapeutic response. Other uses of polyalkylene oxide conjugated sFv proteins are specific targeting of pro-drug activating enzymes to tumor cells by a bispecific molecule with specificity for tumor cells and enzyme. Polyalkylene oxide conjugated sFv may be used for specific delivery of drug to an *in vivo* target, such as a tumor, delivery of radioactive metals for tumor radioimmunodiagnosis or radioimmunotherapy (Goldenberg, D.M., *Am. J. Med.* 94:297 (1993)), nonradioactive metals in applications such as with boron/uranium-neutron capture therapy (Ranadive, G.N., *et al.*, *Nucl. Med. Biol.* 20:1 (1993); Barth, R.F., *et al.*, *Bioconjug. Chem.* 5:58 (1994)), and nuclear magnetic resonance imaging (Sieving, P.F., *et al.*, *Bioconjug. Chem.* 1:65 (1990)). This list is illustrative only.

The invention also extends to uses for the polyalkylene oxide conjugated sFv proteins in purification and biosensors. Affinity purification is made possible by affixing the polyalkylene oxide conjugated sFv protein to a support, with the antigen-binding sites exposed to and in contact with the ligand molecule to be separated, and thus purified. Biosensors generate a detectable signal upon binding of a specific antigen to an antigen-binding molecule, with subsequent processing of the signal. Polyalkylene oxide conjugated sFv proteins, when used as the antigen-binding molecule in biosensors, may change conformation upon binding, thus generating a signal that may be detected.

The invention is also directed to a method of detecting an antigen suspected of being in a sample by contacting the sample with the polyalkylene oxide conjugated sFv that is labeled. A sample may comprise at least one compound, mixture, surface, solution, emulsion, suspension, mixture, cell culture, fermentation culture, cell, tissue, secretion and/or derivative or extract thereof.

Such samples can also include, e.g., animal tissues, such as blood, lymph, cerebrospinal fluid (CNS), bone marrow, gastrointestinal contents, and portions,

cells or internal and external secretions of skin, heart, lung and respiratory system, liver, spleen, kidney, pancreas, gall bladder, gastrointestinal tract, smooth, skeletal or cardiac muscle, circulatory system, reproductive organs, auditory system, the autonomic and central nervous system, and extracts or cell cultures thereof. Such
5 samples can be measured using methods of the present invention *in vitro*, *in vivo* and *in situ*.

Such samples can also include environmental samples such as earth, air or water samples, as well as industrial or commercial samples such as compounds, mixtures, surfaces, aqueous chemical solutions, emulsions, suspensions or
10 mixtures.

Additionally, samples that can be used in methods of the present invention include cell culture and fermentation media used for growth of prokaryotic or eukaryotic cells and/or tissues, such as bacteria, yeast, mammalian cells, plant cells and insect cells.

15 Essentially all of the uses for which monoclonal or polyclonal antibodies, or fragments thereof, have been envisioned by the prior art, can be addressed by the polyalkylene oxide conjugated sFv proteins of the present invention. These uses include detectably-labeled forms of the polyalkylene oxide conjugated sFv protein. Types of labels are well-known to those of ordinary skill in the art. They
20 include radiolabeling, chemiluminescent labeling, fluorochromic labeling, and chromophoric labeling. Other uses include imaging the internal structure of an animal (including a human) by administering an effective amount of a labeled form of the polyalkylene oxide conjugated sFv protein and measuring detectable radiation associated with the animal. They also include improved immunoassays,
25 including sandwich immunoassay, competitive immunoassay, and other immunoassays wherein the labeled antibody can be replaced by the PEGylated sFv protein of this invention. See, e.g., Kohler *et al.*, *Nature* 256:495 (1975); Kohler *et al.*, *Eur. J. Immunol.* 6:511 (1976); Kohler *et al.*, *Eur. J. Immunol.* 6:292 (1976); Hammerling *et al.*, In: *Monoclonal Antibodies and T-Cell Hybridomas*,

pp. 563-681, Elsevier, N (1981); Sambrook *et al.*, *Molecular Cloning - A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory (1989).

Administration

Administration of polyalkylene oxide conjugated sFv conjugates of the
5 invention for *in vivo* diagnostic and therapeutic applications will be by analogous
methods to sFv where the diagnostic or therapeutic principle is directly linked to
the sFv or a loaded carrier is linked by random binding to amine or carboxyl
groups on amino acid residues of the sFv in a non-site-specific manner.

Conjugates of the present invention (immunoconjugates) can be formulated
10 according to known methods to prepare pharmaceutically useful compositions,
such as by admixture with a pharmaceutically acceptable carrier vehicle. Suitable
vehicles and their formulation are described, for example, in *Remington's Pharma-
ceutical Sciences*, 18th ed., Osol, A., ed., Mack, Easton PA (1990). In order to
form a pharmaceutically acceptable composition suitable for effective
15 administration, such compositions will contain a therapeutically effective amount
of the immunoconjugate, either alone, or with a suitable amount of carrier vehicle.

Additional pharmaceutical methods may be employed to control the
duration of action. Controlled release preparations may be achieved by the use of
polymers to complex or absorb the immunoconjugate of the present invention.
20 The controlled delivery may be exercised by selecting appropriate macromolecules
(for example, polyesters, polyamino acids, polyvinyl pyrrolidone, ethylene-
vinylacetate, methylcellulose, carboxymethylcellulose, or protamine sulfate). The
rate of drug release may also be controlled by altering the concentration of such
macromolecules. Another possible method for controlling the duration of action
25 comprises incorporating the therapeutic agents into particles of a polymeric
substance such as polyesters, polyamino acids, hydrogels, poly(lactic acid) or
ethylene vinylacetate copolymers. Alternatively, it is possible to entrap the
immunoconjugate of the invention in microcapsules prepared, for example, by

coacervation techniques or by interfacial polymerization, for example, by the use of hydroxymethylcellulose or gelatin-microcapsules or poly(methylmethacrylate) microcapsules, respectively, or in a colloid drug delivery system, for example, liposomes, albumin microspheres, microemulsions, nanoparticles, nanocapsules, or in macroemulsions. Such teachings are disclosed in *Remington's Pharmaceutical Sciences*, 16th ed., Osol, A., ed., Mack, Easton PA (1990).

The immunoconjugate may be provided to a patient by means well known in the art. Such means of introduction include oral means, intranasal means, subcutaneous means, intramuscular means, intravenous means, intra-arterial means, or parenteral means. Intravenous, intraarterial or intrapleural administration is normally used for lung, breast, and leukemic tumors. Intraperitoneal administration is advised for ovarian tumors. Intrathecal administration is advised for brain tumors and leukemia. Subcutaneous administration is advised for Hodgkin's disease, lymphoma and breast carcinoma. Catheter perfusion is useful for metastatic lung, breast or germ cell carcinomas of the liver. Intralesional administration is useful for lung and breast lesions.

For therapeutic or diagnostic applications, compositions according to the invention may be administered parenterally in combination with conventional injectable liquid carriers such as sterile pyrogen-free water, sterile peroxide-free ethyl oleate, dehydrated alcohol, or propylene glycol. Conventional pharmaceutical adjuvants for injection solution such as stabilizing agent, solubilizing agents and buffers, such as ethanol, complex forming agents such as ethylene diamine tetraacetic acid, tartrate and citrate buffers, and high-molecular weight polymers such as polyethylene oxide for viscosity regulation may be added. Such compositions may be injected intramuscularly, intraperitoneally, or intravenously.

Further non-limiting examples of carriers and diluents include albumin and/or other plasma protein components such as low density lipoproteins, high density lipoproteins and the lipids with which these serum proteins are associated. These lipids include phosphatidyl choline, phosphatidyl serine, phosphatidyl

ethanolamine and neutral lipids such as triglycerides. Lipid carriers also include, without limitation, tocopherol.

At least one polyalkylene oxide conjugated sFv linked to a therapeutic agent according to the invention may be administered by any means that achieve
5 their intended purpose, for example, to treat various pathologies, such as cell inflammatory, allergy, tissue damage or other related pathologies.

A typical regimen for preventing, suppressing, or treating various pathologies comprises administration of an effective amount of an sFv conjugate, administered over a period of one or several days, up to and including between
10 one week and about 24 months.

It is understood that the dosage of the present invention administered *in vivo* or *in vitro* will be dependent upon the age, sex, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired. The ranges of effective doses provided below are not
15 intended to limit the invention and represent preferred dose ranges. However, the most preferred dosage will be tailored to the individual subject, as is understood and determinable by one of skill in the art, without undue experimentation. See, e.g., Berkow *et al.*, eds., *Merck Manual*, 16th edition, Merck and Co., Rahway, N.J. (1992); Goodman *et al.*, eds., *Goodman and Gilman's The Pharmacological*
20 *Basis of Therapeutics*, 8th edition, Pergamon Press, Inc., Elmsford, N.Y. (1990); *Avery's Drug Treatment: Principles and Practice of Clinical Pharmacology and Therapeutics*, 3rd edition, ADIS Press, LTD., Williams and Wilkins, Baltimore, MD. (1987), Ebadi, *Pharmacology*, Little, Brown and Co., Boston (1985), Katzung, *Basic and Clinical Pharmacology*, Appleton and Lange, Norwalk, Conn.
25 (1992), which references and references cited therein, are entirely incorporated herein by reference.

The total dose required for each treatment may be administered by multiple doses or in a single dose. Effective amounts of a diagnostic/pharmaceutical compound or composition of the present invention are from about 0.001 μ g to
30 about 100 mg/kg body weight, administered at intervals of 4-72 hours, for a

period of 2 hours to 5 years, or any range or value therein, such as 0.01-1.0, 1.0-10, 10-50 and 50-100 mg/kg, at intervals of 1-4, 6-12, 12-24 and 24-72 hours, for a period of 0.5, 1.0-2.0, 2.0-4.0 and 4.0-7.0 days, or 1, 1-2, 2-4, 4-52 or more weeks, or 1, 2, 3-10, 10-20, 20-60 or more years, or any range or value therein.

5 Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions, which may contain auxiliary agents or excipients which are known in the art. Pharmaceutical compositions such as tablets and capsules can also be prepared according to routine methods. See, e.g., Berker, *supra*, Goodman, *supra*, Avery, *supra* and Ebadi, *supra*, which
10 are entirely incorporated herein by reference, including all references cited therein.

 Pharmaceutical compositions comprising at least one type of sFv conjugate of the invention, or, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 types of sFv conjugates, of the present invention may be contained in an amount effective to achieve its intended purpose. In addition to at least one sFv conjugate, a
15 pharmaceutical composition may contain suitable pharmaceutically acceptable carriers, such as excipients, carriers and/or auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically.

 Pharmaceutical compositions may also include suitable solutions for administration intravenously, subcutaneously, dermally, orally, mucosally or
20 rectally, and contain from about 0.01 to 99 percent, preferably from about 20 to 75 percent of active component (i.e., the sFv) together with the excipient. Pharmaceutical compositions for oral administration include tablets and capsules. Compositions which can be administered rectally include suppositories. See, e.g., Berker, *supra*, Goodman, *supra*, Avery, *supra* and Ebadi, *supra*. Additional lipid
25 and lipoprotein drug delivery systems that may be included herein are described more fully in *Annals N.Y. Acad. Sci.* 507:775-88, 98-103, and 252-271, which disclosure is hereby incorporated by reference.

 The compositions may also be formulated into orally administrable compositions containing one or more physiologically compatible carriers or
30 excipients, and may be solid or liquid in form. These compositions may, if desired,

contain conventional ingredients such as binding agents, for example, syrups, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, such as lactose, mannitol, starch, calcium phosphate, sorbitol, cyclodextran, or methylcellulose; lubricants such as magnesium stearate, high molecular weight
5 polymers such as polyethylene glycols, high molecular weight fatty acids such as stearic acid or silica; disintegrants such as starch; acceptable wetting agents as, for example, sodium lauryl sulfate.

The oral compositions may assume any convenient form, such as tablets, capsules, lozenges, aqueous or oily suspensions, emulsions, or dry products
10 suitable for reconstitution with water or other liquid medium prior to use. The liquid oral forms may, of course, contain flavors, sweeteners, preservatives such as methyl or propyl p-hydroxybenzoates; suspending agents such as sorbitol, glucose or other sugar syrup, methyl, hydroxymethyl, or carboxymethyl celluloses or gelatin; emulsifying agents such as lecithin or sorbitan monooleate or
15 thickening agents. Non-aqueous compositions may also be formulated which comprise edible oils as, for example, fish-liver or vegetable oils. These liquid compositions may conveniently be encapsulated in, for example, gelatin capsules in a unit dosage amount.

The pharmaceutical compositions according to the present invention may
20 also be administered, if appropriate, either topically as an aerosol or, formulated with conventional bases as a cream or ointment.

The pharmaceutical compositions of the present invention can also be administered by incorporating the active ingredient into colloidal carriers, such as liposomes. Liposome technology is well known in the art, having been described
25 by Allison *et al.*, *Nature* 252:252-254 (1974), and Dancy *et al.*, *J. Immunol.* 120:1109-1113 (1978).

Having now generally described this invention, the same will be better understood by reference to certain specific examples, which are included for the purpose of illustration and not intended to be limiting unless otherwise specified.

Examples

Example 1

Preparation of Methoxypoly(ethylene glycol)-succinimidyl carbonate

(SC-PEG)

5 Dissolve 60 g of methoxy poly(ethylene glycol) (MW 5,000) in 200 ml of 3/1 toluene/dichloromethane and treat with a toluene solution of phosgene (30 ml, 57 mmol) overnight. Evaporate the solution to dryness and remove the remainder of the phosgene under vacuum. Redissolve the residue in 150 ml of 2/1
10 toluene/dichloromethane. Treat the resulting solution with 2.1 g (18 mmol) of solid N-hydroxysuccinimide, followed by 1.7 ml (12 mmol) of triethylamine. Allow the solution to stand for three hours and then filter it and evaporate it to dryness. Dissolve the residue in 600 ml of warm (50° C) ethyl acetate, filter the solution, and cool it to facilitate precipitation of the polymer. Collect the product
15 by filtration, then recrystallize from ethyl acetate, and dry under vacuum over P₂O₅.

Example 2

Preparation of CC 49/212 SCA

20 In the production of monovalent or multivalent antigen-binding proteins, the same recombinant *E. coli* production system that was used for prior single-chain antigen-binding protein production was used. See Bird *et al.*, *Science* 242:423 (1988). This production system produced between 2 and 20% of the total *E. coli* protein as single-chain antigen-binding protein. For protein recovery, the frozen cell paste from three 10-liter fermentations (600-900 g) was thawed
25 overnight at 4°C and gently resuspended at 4°C in 50 mM Tris-HCl, 1.0 mM EDTA, 100 mM KCl, 0.1 mM PMSF, pH 8.0 (lysis buffer), using 10 liters of lysis buffer for every kilogram of wet cell paste. When thoroughly resuspended, the chilled mixture was passed three times through a Manton-Gaulin cell homogenizer

to totally lyse the cells. Because the cell homogenizer raised the temperature of the cell lysate to $25 \pm 5^{\circ}\text{C}$, the cell lysate was cooled to $5 \pm 2^{\circ}\text{C}$ with a Lauda/Brinkman chilling coil after each pass. Complete lysis was verified by visual inspection under a microscope.

5 The cell lysate was centrifuged at 24,300g for 30 minutes at 6°C using a Sorvall RC-5B centrifuge. The pellet containing the insoluble single-chain antigen-binding protein was retained, and the supernatant was discarded. The pellet was washed by gently scraping it from the centrifuge bottles and resuspending it in 5 liters of lysis buffer/kg of wet cell paste. The resulting 3.0-
10 to 4.5-liter suspension was again centrifuged at 24,300g for 30 minutes at 6°C , and the supernatant was discarded. This washing of the pellet removes soluble *E. coli* proteins and can be repeated as many as five times. At any time during this washing procedure the material can be stored as a frozen pellet at -20°C . A substantial time saving in the washing steps can be accomplished by utilizing a
15 Pellicon tangential flow apparatus equipped with 0.22- μm microporous filters, in place of centrifugation.

 The washed pellet was solubilized at 4°C in freshly prepared 6 M guanidine hydrochloride, 50 mM Tris-HCl, 10 mM CaCl_2 , 50 mM KCl, pH 8.0 (dissociating buffer), using 9 ml/g of pellet. If necessary, a few quick pulses from a Heat
20 Systems Ultrasonics tissue homogenizer can be used to complete the solubilization. The resulting suspension was centrifuged at 24,300g for 45 minutes at 6°C and the pellet was discarded. The optical density of the supernatant was determined at 280 nm and if the OD_{280} was above 30, additional dissociating buffer was added to obtain an OD_{280} of approximately 25.

25 The supernatant was slowly diluted into cold ($4-7^{\circ}\text{C}$) refolding buffer (50 mM Tris-HCl, 10 mM CaCl_2 , 50 mM KCl, pH 8.0) until a 1:10 dilution was reached (final volume 10 - 20 liters). Re-folding occurs over approximately eighteen hours under these conditions. The best results are obtained when the GuHCl extract is slowly added to the refolding buffer over a two hour period,
30 with gentle mixing. The solution was filtered through a 0.2 μm Millipore Millipak

200. This filtration step may be optionally preceded by a centrifugation step. The filtrate was concentrated to 1 to 2 liters using an Amicon spiral cartridge with 10,000 MWCO cartridge, again at 4°C.

5 The concentrated crude antigen-binding protein sample was dialyzed against Buffer G (60 mM MOPS, 0.5 mM Ca acetate, pH 6.0 - 6.4) until the conductivity was lowered to that of Buffer G. The sample was then loaded on a 21.5 x 250-mm polyaspartic acid PolyCAT A column, manufactured by Poly LC of Columbia, Maryland. If more than 60 mg of protein is loaded on this column, the resolution begins to deteriorate; thus, the concentrated crude sample often
10 must be divided into several PolyCAT A runs. Most antigen-binding proteins have an extinction coefficient of about $2.0 \text{ ml mg}^{-1} \text{ cm}^{-1}$ at 280 nm and this can be used to determine protein concentration. The antigen-binding protein sample was eluted from the PolyCAT A column with a 50-min linear gradient from Buffer G to Buffer H (60 mM MOPS, 20 mM Ca Acetate, pH 7.5 - 8.0). Most of the
15 single-chain proteins elute between 20 and 26 minutes when this gradient is used. This corresponds to an eluting solvent composition of approximately 70% Buffer G and 30% Buffer H. Most of the bivalent antigen-binding proteins elute later than 45 minutes, which correspond to over 90% Buffer H.

Example 3

20 ***Modification of CC 49/212 Single Chain Antigen Binding Molecule
With SC-PEG***

A sample of CC 49/212 single chain antigen binding molecule (MW=27000) dissolved in KPO_4/NaCl buffer (pH 7.2) was obtained as described in Example 2. The protein was found to be pure using SDS-PAGE (sodium
25 dodecyl sulfate-polyacrylamide gel electrophoresis) and size exclusion chromatography. The concentration of the protein was 0.79 mg/ml. It was further concentrated to at least 2 mg/ml using an Amicon concentrator with a 10,000 dalton nominal size cut-off, i.e., anything greater than 10K is retained.

The modification reaction, i.e., the coupling of the SC-PEG to the CC 49/212, was carried out in 50 mM KPO₄, 150 mM NaCl buffer, which was the storage buffer the protein was supplied in. The pH was raised from 7.2 to 7.5. SC-PEG (MW 5,000) was added in a 50x molar excess to protein. At specific
5 time intervals, the coupling reaction was terminated by the addition of a 50x molar excess of glycine and the extent and progress of the coupling reaction was checked as a function of time using both size exclusion chromatography using a DuPont Zorbax 250 column and SDS-PAGE.

Free SC-PEG remaining in the samples was removed by extensive dialysis
10 on an Amicon Centricon 10.

The samples were checked for degree of modification by size exclusion chromatography. After concentrating the samples, the residual amine concentration on the protein was determined by titration with trinitrobenzene sulfonate and the percentage of amine groups that had reacted with the SC-PEG
15 (the "% modification") was calculated from the results.

Dansyl derivatives of native single chain antigen binding molecule (CC 49/212) and hemoglobin, PEG SCA and hemoglobin, and N-acetyl lysine were prepared. These samples were then analyzed for amino acid. The results of this experiment are shown in Table 3.

-64-

Table 3			
Reaction Time (Minutes)	Protein (mg/ml)	% Modification	Molecular Weight ¹ (Kilodaltons)
0	0.74148	0	27
15	1.3569	52	84
30	1.3587	62	156
60	1.2706	63	
80			224
90			247
120	0.78	65	
¹ Molecular weights were determined by size exclusion chromatography after calibration against standards.			


Example 4

Competition ELISA

The CC49 monoclonal antibody was developed by Dr. Jeffrey Schlom's group, Laboratory of Tumor Immunology and Biology, National Cancer Institute. It binds specifically to the pan-carcinoma tumor antigen TAG-72. See Muraro, R. *et al.*, Cancer Research 48: 4588-4596 (1988).

Figure 1 is a graphical representation of three competition ELISA's in which unlabeled PEG modified CC49/212 single-chain Fv (closed squares), CC49/212 single-chain Fv (open squares), CC49 IgG (open circles), and MOPC-21 IgG (+) competed against a CC49 IgG radiolabeled with ¹²⁵I for binding to the TAG-72 antigen on a human breast melanoma extract. MOPC-21 is a control antibody that does not bind to TAG-72 antigen. In this experiment, 50% competition of ¹²⁵I-CC49 IgG binding required about 200 nM of CC49 IgG, about 550 nM of CC49/212 sFv, and about 3000 nM of PEG modified CC49/212 sFv.


Example 5
Preparation of U-PEG-OH



Materials

- Methoxypoly(ethylene glycol) (m-PEG) was obtained from Union Carbide.
- 5 The solvents were obtained from Aldrich Chemical of Milwaukee, Wisconsin. The methoxypoly(ethylene glycol)-N-succinimidyl carbonate (SC-PEG) was prepared as described in U.S. Patent No. 5,122,614, using m-PEG having a molecular weight of about 5,000. Each of the products prepared in Examples 5-10 was confirmed structurally by carbon-13 NMR.
- 10 The branched polymer, U-PEG-OH, was prepared by adding 100 mg (1.1 mmol) of 1,3-diamino-2-propanol to a solution of 10.0 g (2 mmol) of SC-PEG in 50 mL of methylene chloride. The mixture was stirred for 18 hours at room temperature then filtered. Excess solvent was removed by distillation *in vacuo*. The residue was recrystallized from 2-propanol to yield 7.1 g of product (70%
- 15 yield).

Example 6
Preparation of U-PNP-PEG




-66-

The compound of Example 5 was activated with p-nitrophenyl chloroformate. First, 5.0g (0.5 mmol) of U-PEG-OH was azeotropically dried by refluxing in 75 mL of toluene for 2 hours, resulting in the removal of 25 mL of solvent/water. The reaction mixture was cooled to 30°C, followed by the addition
 5 of 120 mg (0.6 mmol) of p-nitrophenyl chloroformate and 50 mg (0.6 mmol) of pyridine. The resulting mixture was stirred for two hours at 45°C, followed by stirring overnight at room temperature.

The reaction mixture was then filtered through CELITE™, followed by removal of the solvent from the filtrate by distillation *in vacuo*. The residue was
 10 recrystallized from 2-propanol to yield 4.2 g (81% yield) of the product.


Example 7
Preparation of US-PEG



In this example, the U-PNP-PEG of Example 6 was reacted with
 15 N-hydroxysuccinimide to form the succinimidyl carbonate ester of U-PEG. A solution containing 5.0 g (0.5 mmol) of the U-PNP-PEG, 0.6 g (5 mmol) of N-hydroxysuccinimide and 0.13 g (1 mmol) of diisopropylethylamine in 40 ml of methylene chloride was refluxed for 18 hours. The solvent was then removed by distillation *in vacuo*, and the residue was recrystallized from 2-propanol to yield
 20 4.2 g of the succinimidyl carbonate ester (82% yield).

-67-

Example 8
Preparation of XU-PEG-OH




This branched polymer was prepared by reacting the U-PNP-PEG of Example 6 with 2-(2-aminoethoxy) ethanol (i.e., the amino alcohol was reacted
5 with the p-nitrophenyl carbonate). The recrystallized product yield was 86%.

Example 9
Preparation of XU-PNP-PEG

The compound of Example 8 was functionalized with p-nitrophenyl
10 carbonate as in Example 6. The recrystallized product yield was 83%.

Example 10
Preparation of XUS-PEG



In this example, the succinimidyl carbonate derivative of compound
15 prepared in Example 8 was prepared according to the process described in

-68-

Example 7, by reacting N-hydroxysuccinimide with the p-nitrophenyl carbonate derivative of Example 9. The recovered product yield was 84%.

Example 11

Modification of CC49/218 with SC-PEG or XUS-PEG

5

A sample containing CC49/218 was desalted on a PD-10 column in a buffer consisting of 0.1M sodium phosphate, pH 8.0. An equimolar amount of SC-PEG or XUS-PEG was added and the reactions were incubated at 4°C, overnight. The reactions were quenched with an excess of glycine. The modified
10 CC49/218 conjugates were GPC purified and then concentrated in a centricon-10.

The yield based on GPC integration was about 50% for SC-PEG modified CC49/218 and about 40% for XUS-PEG modified CC49/218. The GPC profiles were almost identical to those obtained when the reaction was performed at pH9.0, room temperature. SDS-PAGE revealed that the appropriate derivatives
15 had been made.

Example 12

Competition ELISA

The assay was performed as in Example 4 above using SC-PEG modified CC49/218 and XUS-PEG modified CC49/218 along with the appropriate controls.
20 The results are shown in Figure 4 and in Table 4 below.

5

10

Sample	50% Inhibition (nM)
CC49 IgG	10
CC49/218 sFv	80
SC Unreacted #049301	300
SC Reacted #049304	650
XUS Unreacted #049302	280
XUS Reacted #049303	320

Thus, the affinity of the SC-PEG modified CC49-SCA was within about 8 to 10 fold of the native CC49-SCA and the affinity the XUS-PEG modified CC49-SCA was within about 4 to 5 fold of the native CC49-SCA.

15

Samples #049304 and #049303 were PEG modified, while samples #04901 and #049302 were unmodified CC49/218 isolated from the reaction mixtures.

Example 13

Pharmacokinetics of Plasma Retention of sFv and PEG-sFv

20

Sixty µg of CC49/218 sFv protein or 60 µg of PEG-modified sFv protein were injected intravenously at time 0 into ICR (CD-1) female mice (Harlan - 25g, 7-8 weeks old). Mice were bled at the time points indicated in Figure 5. The percent retention in plasma was quantitated by ELISA methods. For the PEG-modified conjugate, CC49/218 sFv was conjugated to SC-PEG of molecular mass 20,000 (the protocol is described in U.S. Patent 5,122,614, which disclosure is incorporated herein by reference). The average PEG:sFv molar ratio in the tested PEG-sFv conjugate was approximately 1:1.

25

Example 14

PEGylated Multimer Single Chain Antibodies

Using bifunctional PEG, dimers and trimers of CC49-SCA have been made. CC49-SCA was modified with bifunctional PEG as follows: 2 ml of CC49-SCA in phosphate buffered saline (25 mM sodium phosphate, pH 7.3, 0.15 M NaCl) at a concentration of 1.5 mg/ml was modified as follows. Bifunctional PEG (polyethylene glycol with reactive SC at both terminal ends), 1.887 mg (powder) was dissolved in 0.1 ml of MOPS (3-[N-morpholino]propane)-sulfonic acid) buffered at pH 7.3. This PEG solution was added to CC49-SCA solution within 10 seconds of dissolution. The mixture was then stirred at 24° C for 1 hour. At the end of the reaction, solid Guanidine HCl was added to the reaction mixture to a final concentration of 6 M in order to break up non covalently associated CC49-SCA. This material was immediately applied to a size exclusion column (2 cm X 60 cm, Superdex-75) previously equilibrated in the buffer composed of 60 mM Guanidine HCl in 50 mM Tris pH 7.3, 1mM CaCl₂, 0.1 mM PMSF (phenyl methyl sulfonyl flouride), and 50 mM KCl. Multimers of different molecular weights were then fractionated from the column.

These multimers are CC49-SCA separated by a long stretch of PEG (5000 MW, about 226 carbons in length) and were freshly refolded and thus, are not believed to arise from aggregation. It is less likely that there was diabody or multivalent SCA formed as a result of self association of the CC49-SCA. Further evidence was demonstrated by the fact that there was negligible native CC49-SCA in the denaturing SDS-PAGE profile.

The SDS-PAGE electrophoresis patterns for dimeric CC49-SCA, trimeric CC49-SCA, PEG-CC49-SCA and native CC49-SCA under reducing conditions are shown in Figure 6.

The multimers were assayed for binding affinity using the following assay which was modified method described in B Friquet *et al. J. of Immunology Methods*, 77:305-319 (1985). Briefly, various amounts of a given modified single

-71-

chain antibody were mixed with various amounts of the antigen mucin in PBS (phosphate buffered saline). The binding reaction was allowed to reach equilibration for at least 24 hours at 4°C. At the end of the incubation, the unbound CC49-SCA fractions were assayed by ELISA, while the bound fractions were washed away. The total amount of free CC49-SCA was determined by the ELISA using the CC49-SCA sample pre-incubated in the absence of the antigen mucin. The bound antibody was determined by subtraction of the free (unbound) amount from the total amount as determined by ELISA. Since the total amount of the CC49-SCA was known, it was also used as its own standard curve. Note that each type of CC49-SCA had its own reference control. In essence, the protocol was measuring the unbound amount of a particular version of PEG-CC49-SCA as a result of the binding to the antigen. Although PEG may affect the detection reagent in ELISA, this was well contained in the standard references. Therefore, the amount measured was not due to the difference of various PEG on the various versions of PEG-CC49-SCA.

In sum, in this study, increasing concentrations of the CC49-SCA protein were allowed to bind to a fixed amount of the antigen. The amount that binds to 50% of the maximal level is a good indication of the affinity. This data indicate that the affinity of PEG-Di-CC49-SCA and PEG-Tri-CC49-SCA are very similar to that of native CC49-SCA. However, the PEG-modified CC49-SCA monomer had much lower affinity. The binding data are shown in Figure 7.

Example 15

Pharmacokinetics of PEG-CC49-SCA

The pharmacokinetic study of various forms of PEG-CC49-SCA was performed as in Example 13, above.

The data obtained from this study indicate the following:

There was a trend toward longer circulation half-lives as the size of the PEG increases. As more PEG is attached to the protein, the circulation time increases. However, attachment of a few strands of high molecular weight PEG gives a better increase in circulation half-life than multiple stands of lower
5 molecular weight PEG.

The circulation half-life of CC49-SCA-U-PEG, made with the US-PEG prepared in Example 7, was about the same as that of CC49-SCA-PEG-12000. Therefore, the shape of the PEG does not affect the circulation half-life.

Whether the linker is an SC-bond, Flan-bond, hydrazine bond, or TPC,
10 there was no significant change in the circulation half-life. Therefore, the chemical bonds of the linkers, if not releasable, do not affect the circulation half-life. In addition, the PEG remains attached to the protein during the observable time.

The circulation half-life was shortened by carbohydrate. However, if PEG was attached to the carbohydrate, it increases the circulation by about 10 fold.
15 This was not better, however, than attaching an equivalent number of PEG at other sites on CC49-SCA.

The results of the study are shown in the table below:

Pharmacokinetic Data of SCA and PEG-SCA

Native CC49 or PEG used			# PEG per SCA	Circulatory half-life (hours)		Area Under the Curve ($\mu\text{g/ml}\cdot\text{hour}$)		Mean Residence Time (hours)	
MW	Linker	Label		Mean	Std. Error	Mean	Std. Error	Mean	Std. Error
Native CC49/218			0	0.69	0.41	54	27	1	0.59
2,000	SC		2.1	3.41	0.38	135.91	13.02	4.92	0.55
3,400	NHS	biotin	2	2.52	0.45	95.03	14.48	3.64	0.65
5,000	Flan		1	1.81	0.44	67.41	13.98	2.61	0.63
5,000	SC		1.7	4.15	1.08	147.96	32.7	5.98	1.56
5,000	TPC		1.4	3.48	0.95	120.56	28.05	5.02	1.37
5,000	Hz		4.8	9.81	1.61	395.33	56.32	14.15	2.32
10,000	U	biotin	1.5	10.42	1.36	415.86	47.21	15.03	1.96
12,000	Flan		1.2	13.98	1.76	583.19	64.47	20.17	2.53
12,000	SC		1.3	13.27	1.5	586.05	58.08	19.15	2.16
20,000	SC		1	12.86	0.73	1,111	56	18.55	1.05
Native gCC49/3			0	0.39	0.01	36	1	0.56	0.02
5,000	Hz		4.2	3.84	0.46	305	31	5.54	0.66

NOTES:

- A. The PK modeling was following a one-compartment, i.v.-bolus model.
- B. The observed vs. predicted correlations are over 95%.
- C. The circulatory half-life is the time for the drug concentration in the serum to reduce to one half after equilibration is reached (about 2 min.)
- D. AUC = area under the curve, is the integral of the drug blood level over time from zero to last measurement, and is a measure of quantity of drug absorbed and in the body.
- E. Mean residence time is the average amount of time a drug remains in the compartment.
- F. The code for linkers are:
- | | | | |
|----------------------------------|-----------|-------|---------------------------|
| SC = SuccinimidyI | Carbonate | NHS = | N-hydroxy succinimide |
| Hz = Hydrazide | | TPC = | trichlorophenyl carbonate |
| Flan = Thiazolidine thione ester | | U = | SC in the middle of PEG |
- The Flan-bond PEG was prepared as taught in U.S. Patent 5,405,877.

Example 16
Competition Binding Assay

A competition binding assay of biotinylated CC49-SCA with various PEGylated CC49-SCA proteins was performed using an ELISA of the biotinylated CC49-SCA as detected by horseradish-peroxidase conjugated with streptoavidin (SAV-HRP). In the assay, the biotinylated CC49-SCA and PEG-CC49-SCA sample were mixed at various ratios for competition of binding to the antigen mucin on a surface. The amount of biotinylated-CC49-SCA bound to antigen was then measured by SAV-HRP, which would not detect the PEG-modified CC49-SCA. The reduction of binding of biotinylated CC49-SCA due to competition of PEG-CC49-SCA is a reflection of the relative affinity of the two forms of CC49-SCA for the antigen. The level of PEG-CC49-SCA that caused reduction of biotin-CC49-SCA to half of its maximum binding level (IC50) was used in the Cheng-Prusoff formula for determination of the affinity constant K_d as follows: Assuming the biotin-CC49-SCA level is $[s]$ and its affinity is known as K_s , then the affinity for the PEG-CC49-SCA is $K_d = IC_{50}/(1 + [s]/K_s)$. Note that PEG-CC49-SCA was not directly measured and that PEG had no effect on biotin binding because it was on a different molecule. The estimated K_d was then expressed as a percentage of the control (CC49-SCA).

The affinity ranking obtained was as follows:

$CC49 = SC2 > GC = Bio.CC49 = CC12 > C20 = F5 > HZ > PG$

where SC2 is PEG SC2000-CC49-SCA; GC is glyco-CC49-SCA; Bio.CC49-SCA is the biotinylated CC49-SCA; CC12 is the PEG-SC12,000-CC49-SCA; F5 is PEG-Flan-5000-CC49-SCA; HZ is CC49-SCA highly PEGylated on the carboxyl groups with MW 5000 hydrazine-PEG; PG is PEG-glyco-CC49-SCA and is highly PEGylated on the carbohydrate with MW 5000 hydrazine PEG; and C20 is PEG-SC20000-CC49-SCA. The affinity for all the PEG-modified CC49-SCAs shown in Figure 8, were within about two fold of the native CC49-SCA. Taken together with the data in Example 15 (see, the table), the results indicate that

-75-

when the SCA is modified with a lower number of PEG molecules, the resulting affinity of the SCA is better than when the SCA is modified with a higher number of PEG molecules.

5 Although the foregoing refers to particular preferred embodiments, it will be understood that the present invention is not so limited. It will occur to those skilled in the art that various modifications may be made to the disclosed embodiments and that such modifications are intended to be within the scope of the present invention.

-76-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: ENZON, INC.
20 KINGSBRIDGE ROAD
PISCATAWAY, NJ 08854-3963
UNITED STATES OF AMERICA
- APPLICANTS/INVENTORS: WHITLOW, MARC
SHORR, ROBERT G.L.
FILPULA, DAVID R.
LEE, LIHSYNG S.
- (ii) TITLE OF INVENTION: POLYALKYLENE OXIDE-MODIFIED SINGLE CHAIN
POLYPEPTIDES
- (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.
 - (B) STREET: 1100 NEW YORK AVENUE, SUITE 600
 - (C) CITY: WASHINGTON
 - (D) STATE: DC
 - (E) COUNTRY: USA
 - (F) ZIP: 20005
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: To be assigned
 - (B) FILING DATE: Herewith
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 60/044,449
 - (B) FILING DATE: 30-APR-1997
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 60/050,472
 - (B) FILING DATE: 23-JUN-1997
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 60/063,074
 - (B) FILING DATE: 27-OCT-1997
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 60/067,341
 - (B) FILING DATE: 02-DEC-1997
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: JORGE A. GOLDSTEIN
 - (B) REGISTRATION NUMBER: 29,021
 - (C) REFERENCE/DOCKET NUMBER: 0977.184PC02
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 202-371-2600

-77-

(B) TELEFAX: 202-371-2540

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 749 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..738

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GAC	GTC	GTG	ATG	TCA	CAG	TCT	CCA	TCC	TCC	CTA	CCT	GTG	TCA	GTT	GGC	48
Asp	Val	Val	Met	Ser	Gln	Ser	Pro	Ser	Ser	Leu	Pro	Val	Ser	Val	Gly	
1				5					10					15		
GAG	AAG	GTT	ACT	TTG	AGC	TGC	AAG	TCC	AGT	CAG	AGC	CTT	TTA	TAT	AGT	96
Glu	Lys	Val	Thr	Leu	Ser	Cys	Lys	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser	
			20					25					30			
GGT	AAT	CAA	AAG	AAC	TAC	TTG	GCC	TGG	TAC	CAG	CAG	AAA	CCA	GGG	CAG	144
Gly	Asn	Gln	Lys	Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	
		35					40					45				
TCT	CCT	AAA	CTG	CTG	ATT	TAC	TGG	GCA	TCC	GCT	AGG	GAA	TCT	GGG	GTC	192
Ser	Pro	Lys	Leu	Leu	Ile	Tyr	Trp	Ala	Ser	Ala	Arg	Glu	Ser	Gly	Val	
		50				55					60					
CCT	GAT	CGC	TTC	ACA	GGC	AGT	GGA	TCT	GGG	ACA	GAT	TTC	ACT	CTC	TCC	240
Pro	Asp	Arg	Phe	Thr	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Ser	
65					70				75					80		
ATC	AGC	TGT	GTG	AAG	ACT	GAA	GAC	CTG	GCA	GTT	TAT	TAC	TGT	CAG	CAG	288
Ile	Ser	Cys	Val	Lys	Thr	Glu	Asp	Leu	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	
				85				90						95		
TAT	TAT	AGC	TAT	CCC	CTC	ACG	TTC	GGT	GCT	GGG	ACC	AAG	CTT	GTG	CTG	336
Tyr	Tyr	Ser	Tyr	Pro	Leu	Thr	Phe	Gly	Ala	Gly	Thr	Lys	Leu	Val	Leu	
			100					105					110			
AAA	GGC	TCT	TGT	TCC	GGT	AGC	GGC	AAA	CCC	GGG	AGT	GGT	GAA	GGT	AGC	384
Lys	Gly	Ser	Cys	Ser	Gly	Ser	Gly	Lys	Pro	Gly	Ser	Gly	Glu	Gly	Ser	
		115					120					125				
ACT	AAA	GGT	CAG	GTT	CAG	CTG	CAG	CAG	TCT	GAC	GCT	GAG	TTG	GTG	AAA	432
Thr	Lys	Gly	Gln	Val	Gln	Leu	Gln	Gln	Ser	Asp	Ala	Glu	Leu	Val	Lys	
		130				135					140					
CCT	GGG	GCT	TCA	GTG	AAG	ATT	TCC	TGC	AAG	GCT	TCT	GGC	TAC	ACC	TTC	480
Pro	Gly	Ala	Ser	Val	Lys	Ile	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	
145					150				155					160		
ACT	GAC	CAT	GCA	ATT	CAC	TGG	GTG	AAA	CAG	AAC	CCT	GAA	CAG	GGC	CTG	528

-78-

Thr	Asp	His	Ala	Ile	His	Trp	Val	Lys	Gln	Asn	Pro	Glu	Gln	Gly	Leu		
				165					170					175			
GAA	TGG	ATT	GGA	TAT	TTT	TCT	CCC	GGA	AAT	GAT	GAT	TTT	AAA	TAC	AAT		576
Glu	Trp	Ile	Gly	Tyr	Phe	Ser	Pro	Gly	Asn	Asp	Asp	Phe	Lys	Tyr	Asn		
			180					185					190				
GAG	AGG	TTC	AAG	GGC	AAG	GCC	ACA	CTG	ACT	GCA	GAC	AAA	TCC	TCC	AGC		624
Glu	Arg	Phe	Lys	Gly	Lys	Ala	Thr	Leu	Thr	Ala	Asp	Lys	Ser	Ser	Ser		
		195					200					205					
ACT	GCC	TAC	GTG	CAG	CTC	AAC	TGC	CTG	ACA	TCT	GAG	GAT	TCT	GCA	GTG		672
Thr	Ala	Tyr	Val	Gln	Leu	Asn	Cys	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val		
	210					215					220						
TAT	TTC	TGT	ACA	AGA	TCC	CTG	AAT	ATG	GCC	TAC	TGG	GGT	CAA	GGA	ACC		720
Tyr	Phe	Cys	Thr	Arg	Ser	Leu	Asn	Met	Ala	Tyr	Trp	Gly	Gln	Gly	Thr		
225					230					235					240		
TCA	GTC	ACC	GTC	TCC	TGC	TAATAGGATC	C										749
Ser	Val	Thr	Val	Ser	Cys												
				245													

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 246 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Asp	Val	Val	Met	Ser	Gln	Ser	Pro	Ser	Ser	Leu	Pro	Val	Ser	Val	Gly		
1				5					10					15			
Glu	Lys	Val	Thr	Leu	Ser	Cys	Lys	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser		
		20						25					30				
Gly	Asn	Gln	Lys	Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln		
	35						40					45					
Ser	Pro	Lys	Leu	Leu	Ile	Tyr	Trp	Ala	Ser	Ala	Arg	Glu	Ser	Gly	Val		
	50					55					60						
Pro	Asp	Arg	Phe	Thr	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Ser		
65				70					75					80			
Ile	Ser	Cys	Val	Lys	Thr	Glu	Asp	Leu	Ala	Val	Tyr	Tyr	Cys	Gln	Gln		
			85					90						95			
Tyr	Tyr	Ser	Tyr	Pro	Leu	Thr	Phe	Gly	Ala	Gly	Thr	Lys	Leu	Val	Leu		
		100						105					110				
Lys	Gly	Ser	Cys	Ser	Gly	Ser	Gly	Lys	Pro	Gly	Ser	Gly	Glu	Gly	Ser		
	115						120					125					
Thr	Lys	Gly	Gln	Val	Gln	Leu	Gln	Gln	Ser	Asp	Ala	Glu	Leu	Val	Lys		
	130					135					140						

-79-

Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe
 145 150 155 160

Thr Asp His Ala Ile His Trp Val Lys Gln Asn Pro Glu Gln Gly Leu
 165 170 175

Glu Trp Ile Gly Tyr Phe Ser Pro Gly Asn Asp Asp Phe Lys Tyr Asn
 180 185 190

Glu Arg Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser
 195 200 205

Thr Ala Tyr Val Gln Leu Asn Cys Leu Thr Ser Glu Asp Ser Ala Val
 210 215 220

Tyr Phe Cys Thr Arg Ser Leu Asn Met Ala Tyr Trp Gly Gln Gly Thr
 225 230 235 240

Ser Val Thr Val Ser Cys
 245

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 782 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..771

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAC GTC GTG ATG TCA CAG TCT CCA TCC TCC CTA CCT GTG TCA GTT GGC	48
Asp Val Val Met Ser Gln Ser Pro Ser Ser Leu Pro Val Ser Val Gly	
250 255 260	
GAG AAG GTT ACT TTG AGC TGC AAG TCC AGT CAG AGC CTT TTA TAT AGT	96
Glu Lys Val Thr Leu Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser	
265 270 275	
GGT AAT CAA AAG AAC TAC TTG GCC TGG TAC CAG CAG AAA CCA GGG CAG	144
Gly Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln	
280 285 290	
TCT CCT AAA CTG CTG ATT TAC TGG GCA TCC GCT AGG GAA TCT GGG GTC	192
Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Ala Arg Glu Ser Gly Val	
295 300 305 310	
CCT GAT CGC TTC ACA GGC AGT GGA TCT GGG ACA GAT TTC ACT CTC TCC	240
Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Ser	
315 320 325	
ATC AGC AGT GTG AAG ACT GAA GAC CTG GCA GTT TAT TAC TGT CAG CAG	288
Ile Ser Ser Val Lys Thr Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln	
330 335 340	

-80-

TAT TAT AGC TAT CCC CTC ACG TTC GGT GCT GGG ACC AAG CTT GTG CTG	336
Tyr Tyr Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Val Leu	
345 350 355	
AAA GGC TCT ACT TCC GGT AGC GGC AAA CCC GGG AGT GGT GAA GGT AGC	384
Lys Gly Ser Thr Ser Gly Ser Gly Lys Pro Gly Ser Gly Glu Gly Ser	
360 365 370	
ACT AAA GGT CAG GTT CAG CTG CAG CAG TCT GAC GCT GAG TTG GTG AAA	432
Thr Lys Gly Gln Val Gln Leu Gln Gln Ser Asp Ala Glu Leu Val Lys	
375 380 385 390	
CCT GGG GCT TCA GTG AAG ATT TCC TGC AAG GCT TCT GGC TAC ACC TTC	480
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
395 400 405	
ACT GAC CAT GCA ATT CAC TGG GTG AAA CAG AAC CCT GAA CAG GGC CTG	528
Thr Asp His Ala Ile His Trp Val Lys Gln Asn Pro Glu Gln Gly Leu	
410 415 420	
GAA TGG ATT GGA TAT TTT TCT CCC GGA AAT GAT GAT TTT AAA TAC AAT	576
Glu Trp Ile Gly Tyr Phe Ser Pro Gly Asn Asp Asp Phe Lys Tyr Asn	
425 430 435	
GAG AGG TTC AAG GGC AAG GCC ACA CTG ACT GCA GAC AAA TCC TCC AGC	624
Glu Arg Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser	
440 445 450	
ACT GCC TAC GTG CAG CTC AAC AGC CTG ACA TCT GAG GAT TCT GCA GTG	672
Thr Ala Tyr Val Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val	
455 460 465 470	
TAT TTC TGT ACA AGA TCC CTG AAT ATG GCC TAC TGG GGT CAA GGA ACC	720
Tyr Phe Cys Thr Arg Ser Leu Asn Met Ala Tyr Trp Gly Gln Gly Thr	
475 480 485	
TCG GTC ACC GTC TCC AAA AAG AAG AAA AAA AAG AAA AAG GTC ACC GTC	768
Ser Val Thr Val Ser Lys Lys Lys Lys Lys Lys Lys Lys Val Thr Val	
490 495 500	
TCC TAATAGGATC C	782
Ser	

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 257 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Asp	Val	Val	Met	Ser	Gln	Ser	Pro	Ser	Ser	Leu	Pro	Val	Ser	Val	Gly
1				5				10						15	
Glu	Lys	Val	Thr	Leu	Ser	Cys	Lys	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser
			20					25					30		

-81-

Gly Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45
 Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Ala Arg Glu Ser Gly Val
 50 55 60
 Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Ser
 65 70 75 80
 Ile Ser Ser Val Lys Thr Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln
 85 90 95
 Tyr Tyr Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Val Leu
 100 105 110
 Lys Gly Ser Thr Ser Gly Ser Gly Lys Pro Gly Ser Gly Glu Gly Ser
 115 120 125
 Thr Lys Gly Gln Val Gln Leu Gln Gln Ser Asp Ala Glu Leu Val Lys
 130 135 140
 Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe
 145 150 155 160
 Thr Asp His Ala Ile His Trp Val Lys Gln Asn Pro Glu Gln Gly Leu
 165 170 175
 Glu Trp Ile Gly Tyr Phe Ser Pro Gly Asn Asp Asp Phe Lys Tyr Asn
 180 185 190
 Glu Arg Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser
 195 200 205
 Thr Ala Tyr Val Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val
 210 215 220
 Tyr Phe Cys Thr Arg Ser Leu Asn Met Ala Tyr Trp Gly Gln Gly Thr
 225 230 235 240
 Ser Val Thr Val Ser Lys Lys Lys Lys Lys Lys Lys Lys Val Thr Val
 245 250 255

Ser

Claims

What is claimed is:

1. A single-chain antigen-binding polypeptide-polyalkylene oxide conjugate, comprising a single-chain antigen-binding polypeptide comprising:
 - (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;
 - (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and
 - (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,wherein said single-chain antigen-binding polypeptide is conjugated to polyalkylene oxide and
wherein said single-chain antigen-binding polypeptide-polyalkylene oxide conjugate has an antigen binding affinity within a range of about one-fold to about ten-fold of the antigen binding affinity of said single-chain antigen-binding polypeptide in its unconjugated form.
2. A single-chain antigen-binding polypeptide-polyalkylene oxide conjugate, comprising a single-chain antigen-binding polypeptide comprising:
 - (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;
 - (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and
 - (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,

wherein said single-chain antigen-binding polypeptide is conjugated to polyalkylene oxide and

wherein said single-chain antigen-binding polypeptide-polyalkylene oxide conjugate has an antigen binding affinity within about ten-fold of the antigen binding affinity of said single-chain antigen-binding polypeptide in its unconjugated form.

3. A single-chain antigen-binding polypeptide-polyalkylene oxide conjugate, comprising a single-chain antigen-binding polypeptide comprising:

- (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

- (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

- (c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,

wherein said single-chain antigen-binding polypeptide is conjugated to polyalkylene oxide and

wherein said single-chain antigen-binding polypeptide-polyalkylene oxide conjugate has an antigen binding affinity within about five-fold of the antigen binding affinity of said single-chain antigen-binding polypeptide in its unconjugated form.

4. A single-chain antigen-binding polypeptide-polyalkylene oxide conjugate, comprising a single-chain antigen-binding polypeptide comprising:

- (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

- (b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

-84-

(c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site, wherein said single-chain antigen-binding polypeptide is conjugated to polyalkylene oxide and

wherein said single-chain antigen-binding polypeptide-polyalkylene oxide conjugate has an antigen binding affinity within about two-fold of the antigen binding affinity of said single-chain antigen-binding polypeptide in its unconjugated form.

5. A single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

(a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) a peptide linker linking said first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,

wherein the single-chain antigen-binding polypeptide has at least one Cys residue which is capable of polyalkylene oxide conjugation, wherein said Cys residue is located at a position selected from the group consisting of:

(i) the amino acid position 11, 12, 13, 14 or 15 of the light chain variable region;

(ii) the amino acid position 77, 78 or 79 of the light chain variable region;

(iii) the amino acid position 11, 12, 13, 14 or 15 of the heavy chain variable region;

(iv) the amino acid position 82B, 82C or 83 of the heavy chain variable region;

(v) any amino acid position of the peptide linker;

(vi) adjacent to the C-terminus of polypeptide (a) or (b); and

(vii) combinations thereof,

wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

6. The single-chain antigen-binding polypeptide of claim 5, wherein said Cys residue capable of polyalkylene oxide conjugation is located at a position selected from the group consisting of:

- (i') the amino acid position 77 of the light chain variable region;
- (ii') the amino acid position 82B of the heavy chain variable region;
- (iii') the amino acid position 3 of the peptide linker;
- (iv') adjacent to the C-terminus of said polypeptide (a) or (b); and
- (v') combinations thereof.

7. The single-chain antigen-binding polypeptide of claim 5, wherein said first polypeptide (a) comprises the antigen binding portion of the variable region of an antibody light chain and said second polypeptide (b) comprises the antigen binding portion of the variable region of an antibody heavy chain.

8. The single-chain antigen-binding polypeptide of claim 5, wherein the C-terminus of said second polypeptide (b) is the native C-terminus.

9. The single-chain antigen-binding polypeptide of claim 5, wherein the C-terminus of said second polypeptide (b) comprises a deletion of one or plurality of amino acid residue(s), such that the remaining N-terminus amino acid residues of the second polypeptide are sufficient for the polyalkylene oxide conjugated single-chain antigen-binding polypeptide to be capable of binding an antigen.

10. The single-chain antigen-binding polypeptide of claim 5, wherein the C-terminus of said second polypeptide (b) comprises an addition of one or plurality of amino acid residue(s), such that the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

11. The single-chain antigen-binding polypeptide of claim 5, wherein said wherein said Cys residue capable of polyalkylene oxide conjugation is attached to a polyalkylene oxide moiety.

12. The polyalkylene oxide conjugated single-chain antigen-binding polypeptide of claim 11, wherein said polyalkylene oxide conjugated single-chain antigen-binding polypeptide is conjugated to one or plurality of peptide, lipid, nucleic acid, drug, toxin, chelator, boron addend or detectable label molecules.

13. The polyalkylene oxide conjugated single-chain antigen-binding polypeptide of claim 11, wherein said polyalkylene oxide conjugated single-chain antigen-binding polypeptide is conjugated to a carrier having one or plurality of peptide, lipid, nucleic acid, drug, toxin, chelator, boron addend or detectable label molecules bound to said carrier.

14. A polynucleotide sequence encoding a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

(a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) a peptide linker linking said first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,

wherein the single-chain antigen-binding polypeptide has at least one Cys residue which is capable of polyalkylene oxide conjugation, wherein said Cys residue is located at a position selected from the group consisting of:

(i) the amino acid position 11, 12, 13, 14 or 15 of said light chain variable region;

-87-

(ii) the amino acid position 77, 78 or 79 of said light chain variable region;

(iii) the amino acid position 11, 12, 13, 14 or 15 of said heavy chain variable region;

(iv) the amino acid position 82B, 82C or 83 of said heavy chain variable region;

(v) any amino acid position of said peptide linker;

(vi) adjacent to the C-terminus of said polypeptide (a) or (b); and

(vii) combinations thereof, and

wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

15. The polynucleotide sequence of claim 14, wherein said Cys residue capable of polyalkylene oxide conjugation is located at a position selected from the group consisting of:

(i') the amino acid position 77 of the light chain variable region;

(ii') the amino acid position 82B of the heavy chain variable region;

(iii') the amino acid position 3 of the peptide linker;

(iv') adjacent to the C-terminus of said polypeptide (a) or (b); and

(v') combinations thereof.

16. A replicable cloning or expression vehicle comprising the polynucleotide sequence of claim 14.

17. The vehicle of claim 16 which is a plasmid.

18. A host cell transformed with the polynucleotide of claim 17.

19. The host cell of claim 18 which is a bacterial cell, a yeast cell or other fungal cell, an insect cell or a mammalian cell line.

20. The host cell of claim 19 which is *Pichia pastoris*.

21. A method of producing a single-chain antigen-binding polypeptide-polyalkylene oxide conjugate, comprising a single-chain antigen-binding polypeptide comprising:

(a) providing a first genetic sequence encoding a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) providing a second genetic sequence encoding a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) linking the first and second genetic sequences (a) and (b) with a third genetic sequence encoding a linker into a fourth genetic sequence encoding a single chain polypeptide having an antigen binding site,

wherein said single-chain antigen-binding polypeptide is capable of polyalkylene oxide conjugation and

wherein said single-chain antigen-binding polypeptide-polyalkylene oxide conjugate retains antigen binding affinity within a range of about one-fold to about ten-fold of the antigen binding affinity of said single-chain antigen-binding polypeptide in its unconjugated form.

22. A method of producing a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

(a) providing a first genetic sequence encoding a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) providing a second genetic sequence encoding a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) linking said first and second genetic sequences (a) and (b) with a third genetic sequence encoding a peptide linker into a fourth genetic sequence encoding a single chain polypeptide having an antigen binding site,

wherein the single-chain antigen-binding polypeptide has at least one Cys residue which is capable of polyalkylene oxide conjugation, wherein said Cys residue is located at a position selected from the group consisting of:

- (i) the amino acid position 11, 12, 13, 14 or 15 of said light chain variable region;
- (ii) the amino acid position 77, 78 or 79 of said light chain variable region;
- (iii) the amino acid position 11, 12, 13, 14 or 15 of said heavy chain variable region;
- (iv) the amino acid position 82B, 82C or 83 of said heavy chain variable region;
- (v) any amino acid position of said peptide linker;
- (vi) adjacent to the C-terminus of said polypeptide (a) or (b); and
- (vii) combinations thereof, and

wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen;

(d) transforming a host cell with said fourth genetic sequence encoding said single-chain antigen-binding polypeptide of step (c); and

(e) expressing said single-chain antigen-binding polypeptide of step (c) in said host, thereby producing a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation.

23. The method of claim 22, wherein said Cys residue capable of polyalkylene oxide conjugation is located at a position selected from the group consisting of:

- (i') the amino acid position 77 of the light chain variable region;
- (ii') the amino acid position 82B of the heavy chain variable region;

-90-

- (iii') the amino acid position 3 of the peptide linker;
- (iv') adjacent to the C-terminus of said polypeptide (a) or (b); and
- (v') combinations thereof.

24. The method of claim 22, said first genetic sequence encoding a first polypeptide (a) comprises the antigen binding portion of the variable region of an antibody light chain and said second genetic sequence encoding a second polypeptide (b) comprises the antigen binding portion of the variable region of an antibody heavy chain.

25. The method of claim 22, wherein the C-terminus of said second polypeptide (b) is the native C-terminus.

26. The method of claim 22, wherein the C-terminus of said second polypeptide (b) comprises a deletion of one or plurality of amino acid residue(s), such that the remaining N-terminus amino acid residues of the second polypeptide are sufficient for the polyalkylene oxide conjugated single-chain antigen-binding polypeptide to be capable of binding an antigen.

27. The method of claim 22, wherein the C-terminus of said second polypeptide (b) comprises an addition of one or plurality of amino acid residue(s), such that the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

28. A multivalent single-chain antigen-binding polypeptide-polyalkylene oxide conjugate, comprising two or more single-chain antigen binding polypeptides, each single-chain antigen-binding polypeptide comprising:

- (a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

-91-

(b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site, wherein said single-chain antigen-binding polypeptide is conjugated to polyalkylene oxide and

wherein said single-chain antigen-binding polypeptide-polyalkylene oxide conjugate retains antigen binding affinity within a range of about one-fold to about ten-fold of the antigen binding affinity of said single-chain antigen-binding polypeptide in its unconjugated form.

29. A multivalent single-chain antigen-binding protein, comprising two or more single-chain antigen-binding polypeptides, each single-chain antigen-binding polypeptide comprising:

(a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) a peptide linker linking said first and second polypeptides wherein one of the two single-chain antigen-binding polypeptide has at least one Cys residue which is capable of polyalkylene oxide conjugation, wherein said Cys residue is located at a position selected from the group consisting of:

(i) the amino acid position 11, 12, 13, 14 or 15 of said light chain variable region;

(ii) the amino acid position 77, 78 or 79 of said light chain variable region;

(iii) the amino acid position 11, 12, 13, 14 or 15 of said heavy chain variable region;

(iv) the amino acid position 82B, 82C or 83 of said heavy chain variable region;

-92-

- (v) any amino acid position of said peptide linker;
- (vi) adjacent to the C-terminus of said polypeptide (a) or (b); and
- (vii) combinations thereof, and

wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

30. The multivalent protein of claim 29, wherein said Cys residue capable of polyalkylene oxide conjugation is located at a position selected from the group consisting of:

- (i') the amino acid position 77 of the light chain variable region;
- (ii') the amino acid position 82B of the heavy chain variable region;
- (iii') the amino acid position 3 of the peptide linker;
- (iv') adjacent to the C-terminus of said polypeptide (a) or (b); and
- (v') combinations thereof.

31. The multivalent protein of claim 29, wherein said first polypeptide (a) comprises the antigen binding portion of the variable region of an antibody light chain and said second polypeptide (b) comprises the antigen binding portion of the variable region of an antibody heavy chain.

32. The multivalent protein of claim 29, wherein the C-terminus of said second polypeptide (b) is the native C-terminus.

33. The multivalent protein of claim 29, wherein the C-terminus of said second polypeptide (b) comprises a deletion of one or plurality of amino acid residue(s), such that the remaining N-terminus amino acid residues of the second polypeptide are sufficient for the polyalkylene oxide conjugated single-chain antigen-binding polypeptide to be capable of binding an antigen.

34. The multivalent protein of claim 29, wherein the C-terminus of said second polypeptide comprises an addition of one or plurality of amino acid residue(s), such that the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

35. The multivalent protein of claim 29, wherein said wherein said Cys residue capable of polyalkylene oxide conjugation is attached to a polyalkylene oxide moiety.

36. The polyalkylene oxide conjugated multivalent protein of claim 35, wherein said polyalkylene oxide conjugated multivalent protein is conjugated to one or plurality of peptide, lipid, nucleic acid, drug, toxin, chelator, boron addend or detectable label molecule(s).

37. A method of detecting an antigen suspected of being in a sample, comprising:

(a) contacting said sample with the polyalkylene oxide conjugated polypeptide or protein of claim 1 or 11, wherein said polyalkylene oxide conjugated polypeptide or protein is conjugated to one or plurality of detectable label molecule(s), or conjugated to a carrier having one or plurality of detectable label molecule(s) bound to said carrier; and

(b) detecting whether said polyalkylene oxide conjugated single-chain antigen-binding polypeptide or protein has bound to said antigen.

38. A method of imaging the internal structure of an animal, comprising administering to said animal an effective amount of the polyalkylene oxide conjugated polypeptide or protein of claim 1 or 11, wherein said polyalkylene oxide conjugated polypeptide or protein is conjugated to one or plurality of detectable label or chelator molecule(s), or conjugated to a carrier having one or

plurality of detectable label or chelator molecule(s) bound to said carrier, and measuring detectable radiation associated with said animal.

39. The method of claim 38, wherein said animal includes a human.

40. A method for treating a targeted disease, comprising administering an effective amount of a composition comprising the polyalkylene oxide conjugated polypeptide or protein of claim 1 or 11 and a pharmaceutically acceptable carrier vehicle, wherein said polyalkylene oxide conjugated polypeptide or protein is conjugated to one or plurality of peptide, lipid, nucleic acid, drug, toxin, boron addend or radioisotope molecule(s), or conjugated to a carrier having one or plurality of peptide, lipid, nucleic acid, drug, toxin, boron addend or radioisotope molecule(s) bound to said carrier.

41. A single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

(a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,

wherein the single-chain antigen-binding polypeptide has at least three consecutive Lys residues which are capable of polyalkylene oxide conjugation and wherein any one of said consecutive Lys residues is located at a position selected from the group consisting of

(i) any amino acid position of the peptide linker;

(ii) adjacent to the C-terminus of the second polypeptide (b); and

(iii) combinations thereof,

wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

42. A polynucleotide sequence encoding a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

(a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site,

wherein the single-chain antigen-binding polypeptide has at least three consecutive Lys residues which are capable of polyalkylene oxide conjugation and wherein any one of said consecutive Lys residues is located at a position selected from the group consisting of

(i) any amino acid position of the peptide linker;

(ii) adjacent to the C-terminus of the second polypeptide (b); and

(iii) combinations thereof,

wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

43. A replicable cloning or expression vehicle comprising the polynucleotide sequence of claim 42.

44. The vehicle of claim 43 which is a plasmid.

45. A host cell transformed with the polynucleotide of claim 42.

46. The host cell of claim 39 which is a bacterial cell, a yeast cell or other fungal cell, an insect cell or a mammalian cell line.

47. The host cell of claim 46 which is *Pichia pastoris*.

48. A method of producing a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

(a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) a peptide linker linking the first and second polypeptides (a) and (b) into a single chain polypeptide having an antigen binding site, wherein the single-chain antigen-binding polypeptide has at least three consecutive Lys residues which are capable of polyalkylene oxide conjugation and wherein any one of said consecutive Lys residues is located at a position selected from the group consisting of

(i) any amino acid position of the peptide linker;

(ii) adjacent to the C-terminus of the second polypeptide (b); and

(iii) combinations thereof,

wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

49. The single-chain antigen-binding polypeptide of claim 41, wherein said wherein said wherein any one of said consecutive Lys residues capable of polyalkylene oxide conjugation is attached to a polyalkylene oxide moiety.

50. The polyalkylene oxide conjugated single-chain antigen-binding polypeptide of claim 49, wherein said polyalkylene oxide conjugated single-chain antigen-binding polypeptide is conjugated to one or plurality of peptide, lipid, nucleic acid, drug, toxin, chelator, boron addend or detectable label molecules.

51. The polyalkylene oxide conjugated single-chain antigen-binding polypeptide of claim 49, wherein said polyalkylene oxide conjugated single-chain antigen-binding polypeptide is conjugated to a carrier having one or plurality of peptide, lipid, nucleic acid, drug, toxin, chelator, boron addend or detectable label molecules bound to said carrier.

52. A method of producing a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation, comprising:

(a) providing a first genetic sequence encoding a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

(b) providing a second genetic sequence encoding a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) linking said first and second genetic sequences (a) and (b) with a third genetic sequence encoding a peptide linker into a fourth genetic sequence encoding a single chain polypeptide having an antigen binding site,

wherein the single-chain antigen-binding polypeptide has at least three consecutive Lys residues which are capable of polyalkylene oxide conjugation and wherein any one of said consecutive Lys residues is located at a position selected from the group consisting of

- (i) any amino acid position of the peptide linker;
- (ii) adjacent to the C-terminus of the second polypeptide (b); and
- (iii) combinations thereof,

wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen;

(d) transforming a host cell with said fourth genetic sequence encoding said single-chain antigen-binding polypeptide of step (c); and

(e) expressing said single-chain antigen-binding polypeptide of step (c) in said host, thereby producing a single-chain antigen-binding polypeptide capable of polyalkylene oxide conjugation.

53. The method of claim 52, said first genetic sequence encoding a first polypeptide (a) comprises the antigen binding portion of the variable region of an antibody light chain and said second genetic sequence encoding a second polypeptide (b) comprises the antigen binding portion of the variable region of an antibody heavy chain.

54. The method of claim 52, wherein the C-terminus of said second polypeptide (b) is the native C-terminus.

55. The method of claim 52, wherein the C-terminus of said second polypeptide (b) comprises a deletion of one or plurality of amino acid residue(s), such that the remaining N-terminus amino acid residues of the second polypeptide are sufficient for the polyalkylene oxide conjugated single-chain antigen-binding polypeptide to be capable of binding an antigen.

56. The method of claim 52, wherein the C-terminus of said second polypeptide (b) comprises an addition of one or plurality of amino acid residue(s), such that the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

57. A multivalent single-chain antigen-binding protein, comprising two or more single-chain antigen-binding polypeptides, each single-chain antigen-binding polypeptide comprising:

(a) a first polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain;

-99-

(b) a second polypeptide comprising the antigen binding portion of the variable region of an antibody heavy or light chain; and

(c) a peptide linker linking said first and second polypeptides wherein one of the two single-chain antigen-binding polypeptides has at least three consecutive Lys residues which are capable of polyalkylene oxide conjugation and wherein any one of said consecutive Lys residues is located at a position selected from the group consisting of

- (i) any amino acid position of the peptide linker;
- (ii) adjacent to the C-terminus of the second polypeptide (b); and
- (iii) combinations thereof,

wherein the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

58. The multivalent protein of claim 57, wherein said first polypeptide (a) comprises the antigen binding portion of the variable region of an antibody light chain and said second polypeptide (b) comprises the antigen binding portion of the variable region of an antibody heavy chain.

59. The multivalent protein of claim 57, wherein the C-terminus of said second polypeptide (b) is the native C-terminus.

60. The multivalent protein of claim 57, wherein the C-terminus of said second polypeptide (b) comprises a deletion of one or plurality of amino acid residue(s), such that the remaining N-terminus amino acid residues of the second polypeptide are sufficient for the polyalkylene oxide conjugated single-chain antigen-binding polypeptide to be capable of binding an antigen.

61. The multivalent protein of claim 57, wherein the C-terminus of said second polypeptide comprises an addition of one or plurality of amino acid

residue(s), such that the polyalkylene oxide conjugated single-chain antigen-binding polypeptide is capable of binding an antigen.

62. The multivalent protein of claim 57, wherein any one of said consecutive Lys residues capable of polyalkylene oxide conjugation is attached to a polyalkylene oxide moiety.

63. The polyalkylene oxide conjugation multivalent protein of claim 62, wherein said polyalkylene oxide conjugated multivalent protein is conjugated to one or plurality of peptide, lipid, nucleic acid, drug, toxin, chelator, boron addend or detectable label molecule(s).


Figure 1

Figure 2

CC49 V_L 12 15
 D V V M S Q S P S S L P V S V
GAC GTC GTG ATG TCA CAG TCT CCA TCC TCC CTA CCT GTG TCA GTT
 Aat II 27C
 G E K V T L S C K S S Q S L L
 GGC GAG AAG GTT ACT TTG AGC TGC AAG TCC AGT CAG AGC CTT TTA
 39
 Y S G N Q K N Y L A W Y Q Q K
TAT AGT GGT AAT CAA AAG AAC TAC TTG GCC TGG TAC CAG CAG AAA
 54
 P G Q S P K L L I Y W A S A R
 CCA GGG CAG TCT CCT AAA CTG CTG ATT TAC TGG GCA TCC GCT AGG
 69
 E S G V P D R F T G S G S G T
GAA TCT GGG GTC CCT GAT CGC TTC ACA GGC AGT GGA TCT GGG ACA
 77 84
 D F T L S I S C V K T E D L A
 GAT TTC ACT CTC TCC ATC AGC TGT GTG AAG ACT GAA GAC CTG GCA
 * 99
 V Y Y C Q Q Y Y S Y P L T F G
 GTT TAT TAC TGT CAG CAG TAT TAT AGC TAT CCC CTC ACG TTC GGT

218 Linker
 A G T K L V L K G S C S G S G
 GCT GGG ACC AAG CTT GTG CTG AAA GGC TCT TGT TCC GGT AGC GGC
 Hind III *

CC49 V_H 4
 K P G S G E G S T K G Q V Q L
 AAA CCC GGG AGT GGT GAA GGT AGC ACT AAA GGT CAG GTT CAG CTG
 Sma I Pvu II

13 19
 Q Q S D A E L V K P G A S V K
 CAG CAG TCT GAC GCT GAG TTG GTG AAA CCT GGG GCT TCA GTG AAG
 34
 I S C K A S G Y T F T D H A I
 ATT TCC TGC AAG GCT TCT GGC TAC ACC TTC ACT GAC CAT GCA ATT
 49
 H W V K Q N P E Q G L E W I G
CAC TGG GTG AAA CAG AAC CCT GAA CAG GGC CTG GAA TGG ATT GGA
 63
 Y F S P G N D D F K Y N E R F
TAT TTT TCT CCC GGA AAT GAT GAT TTT AAA TAC AAT GAG AGG TTC
 78
 K G K A T L T A D K S S S T A
AAG GGC AAG GCC ACA CTG ACT GCA GAC AAA TCC TCC AGC ACT GCC
 82B 90
 Y V Q L N C L T S E D S A V Y
 TAC GTG CAG CTC AAC TGC CTG ACA TCT GAG GAT TCT GCA GTG TAT
 * 107
 F C T R S L N M A Y W G Q G T
 TTC TGT ACA AGA TCC CTG AAT ATG GCC TAC TGG GGT CAA GGA ACC

Figure 2 (con't)

 112
S V T V S C
TCA GTC ACC GTC TCC TGC TAA TAGGATCC
 * Bam HI

Figure legend. DNA sequence and protein sequence of CC49/218 sFv which has four engineered cysteine residues at the positions indicated by the codons underlined and marked by an asterisk. Also highlighted are the CDR sequences (double underlined) and the 218 linker (underlined and labeled). Note that there are also four natural cysteine residues in the protein which are involved in two disulfide bonds. These are not underlined. The four engineered cysteine residues occur independently in four different mutants currently, but may be combined in the exact four-mutant-codon version shown above.

Figure 3

CC49 V_L 12 15
D V V M S Q S P S S L P V S V
GAC GTC GTG ATG TCA CAG TCT CCA TCC TCC CTA CCT GTG TCA GTT
Aat II 27C
G E K V T L S C K S S Q S L L
GGC GAG AAG GTT ACT TTG AGC TGC AAG TCC AGT CAG AGC CTT TTA
39
Y S G N Q K N Y L A W Y Q Q K
TAT AGT GGT AAT CAA AAG AAC TAC TTG GCC TGG TAC CAG CAG AAA
54
P G Q S P K L L I Y W A S A R
CCA GGG CAG TCT CCT AAA CTG CTG ATT TAC TGG GCA TCC GCT AGG
69
E S G V P D R F T G S G S G T
GAA TCT GGG GTC CCT GAT CGC TTC ACA GGC AGT GGA TCT GGG ACA
77 84
D F T L S I S S V K T E D L A
GAT TTC ACT CTC TCC ATC AGC AGT GTG AAG ACT GAA GAC CTG GCA
99
V Y Y C Q Q Y Y S Y P L T F G
GTT TAT TAC TGT CAG CAG TAT TAT AGC TAT CCC CTC ACG TTC GGT
218 Linker
A G T K L V L K G S T S G S G
GCT GGG ACC AAG CTT GTG CTG AAA GGC TCT ACT TCC GGT AGC GGC
Hind III
CC49 V_H 4
K P G S G E G S T K G Q V Q L
AAA CCC GGG AGT GGT GAA GGT AGC ACT AAA GGT CAG GTT CAG CTG
Sma I Pvu II
13 19
Q Q S D A E L V K P G A S V K
CAG CAG TCT GAC GCT GAG TTG GTG AAA CCT GGG GCT TCA GTG AAG
34
I S C K A S G Y T F T D H A I
ATT TCC TGC AAG GCT TCT GGC TAC ACC TTC ACT GAC CAT GCA ATT
49
H W V K Q N P E Q G L E W I G
CAC TGG GTG AAA CAG AAC CCT GAA CAG GGC CTG GAA TGG ATT GGA
63
Y F S P G N D D F K Y N E R F
TAT TTT TCT CCC GGA AAT GAT GAT TTT AAA TAC AAT GAG AGG TTC
78
K G K A T L T A D K S S S T A
AAG GGC AAG GCC ACA CTG ACT GCA GAC AAA TCC TCC AGC ACT GCC
82B 90
Y V Q L N S L T S E D S A V Y
TAC GTG CAG CTC AAC AGC CTG ACA TCT GAG GAT TCT GCA GTG TAT
107
F C T R S L N M A Y W G Q G T
TTC TGT ACA AGA TCC CTG AAT ATG GCC TAC TGG GGT CAA GGA ACC

Figure 3 (con't)

					112														
S	V	T	V	S	K	K	K	K	K	K	K	K	V	T					
<u>TCG</u>	<u>GTC</u>	<u>ACC</u>	<u>GTC</u>	<u>TCC</u>	<u>AAA</u>	<u>AAG</u>	<u>AAG</u>	<u>AAA</u>	<u>AAA</u>	<u>AAG</u>	<u>AAA</u>	<u>AAG</u>	<u>GTC</u>	<u>ACC</u>					
					*	*	*	*	*	*	*	*							
BstEII															BstEII				

V	S																		
<u>GTC</u>	<u>TCC</u>	<u>TAA</u>	<u>TAGGATCC</u>																
BamHI																			


Figure legend. DNA sequence and protein sequence of CC49/218 sFv with engineered oligo-lysine C-terminal tail segment. The eight new lysine residues were genetically engineered at a BstEII site and are shown underlined and marked with asterisks. Also highlighted are the CDR sequences (double underlined); the 218 linker (underlined and labeled); and selected restriction sites.

Figure 4

Competition RIA

CC491-125

Pegylated CC49/218 sFv



Pharmacokinetics of Plasma Retention of sFv and PEG-sFv




Figure 5

Figure 6






Figure 7

Figure 8

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/08654

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 39/385, 39/395; G01N 33/53; C12P 21/06, 21/04; C07K 16/00

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/193.1, 178.1; 435/7.1, 69.6, 69.7, 69.1; 530/387.1, 810, 807

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE, DIALOG, BIOSIS, EMBASE

polyethylene glycol, PEG, single chain antigen binding polypeptide, polyalkylene oxide conjugates

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4,946,778 A (LADNER et al.) 07 August 1990, see entire document.	1-4, 11-13, 21, 28 and 49-51
Y	US 5,122,614 A (ZALIPSKY) 16 June 1992, see entire document.	1-4, 11-13, 21, 28, 37, 49-51, 62 and 63
Y	CUMBER et al. Comparative stabilities in vitro and in vivo of a recombinant mouse antibody FvCys fragment and a bisFvCys conjugate. J. Immunol. 01 July 1992, Vol. 149, No. 1, pages 120-126, see entire document.	11-13 and 28

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 27 JULY 1998	Date of mailing of the international search report 24 AUG 1998
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>S. Devi</i> Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/08654

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LADD et al. Reagents for the preparation of chromophorically labeled polyethylene glycol-protein conjugates. Anal. Biochem. May 1993, Vol. 210, No. 2, pages 258-261, see entire document.	12-13, 37, 49-51, 62 and 63
Y	NATHAN et al. Copolymers of lysine and polyethylene glycol: A new family of functionalized drug carriers. Bioconj. Chem. January 1993, Vol. 4, No. 1, pages 54-62, see entire document.	49-51, 62 and 63
Y	MONFARDINI et al. A branched monomethoxypoly(ethylene glycol) for protein modification. Bioconj. Chem. January 1995, Vol. 6, No. 1, pages 62-69, see entire document.	49-51, 62 and 63
Y	HERSHFIELD et al. Use of site-directed mutagenesis to enhance the epitope-shielding effects of covalent modification of proteins with polyethylene glycol. Proc. Natl. Acad. Sci. USA. August 1991, Vol. 88, No. 16, pages 7185-7189, see entire document.	21
Y	ANAND et al. Bacterial expression and secretion of various single chain Fv genes encoding proteins specific for a Salmonella serotype B O-antigen. J. Biol. Chem. November 1991, Vol. 266, No. 32, pages 21874-21879, see entire document.	21
Y	WHITLOW et al. Multivalent Fvs: characterization of single-chain Fv oligomers and preparation of a bispecific Fv. Protein Eng. August 1994, Vol. 7, No. 8, pages 1017-1026, see entire document.	28, 62 and 63
Y	ZALIPSKY et al. Poly(ethylene glycol)-grafted liposomes with oligopeptide or oligosaccharide ligands appended to the termini of the polymer chains. Bioconj. Chem. March 1997, Vol. 8, No. 2, pages 111-118, see entire document.	12, 13, 50, 51 and 63
Y	ZALIPSKY S. Synthesis of an end-group functionalized polyethylene glycol-lipid conjugate for preparation of polymer-grafted liposomes. Bioconj. Chem. July 1993, Vol. 4, No. 4, pages 296-299, see entire document.	12, 13, 50, 51 and 63
Y	ZALIPSKY et al. Peptide attachment to extremities of liposomal surface grafted PEG chains: Preparation of the long-circulating form of laminin pentapeptide, YIGSR. Bioconj. Chem. November 1995, Vol. 6, No. 6, pages 705-708, see entire document.	12, 13, 50, 51 and 63
Y	SHARP et al. Synthesis and application of a poly(ethylene glycol)-antibody affinity ligand for cell separation in aqueous polymer two-phase systems. Anal. Biochem. April 1986, Vol. 154, No. 1, pages 110-117, see entire document.	37

INTERNATIONAL SEARCH REPORT

international application No.
PCT/US98/08654

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,438,040 A (EKWURIBE et al.) 01 August 1995 (08/01/95), columns 9-11, see entire document.	37
Y	GEORGE et al. Production of a bispecific antibody by linkage of two recombinant single chain Fv molecules. J. Cellular Biochem. March 1991, Suppl. 15E, abstract number N 206, page 127, see entire abstract.	21
A	ZALIPSKY. Functionalized poly(ethylene glycol) for preparation of biologically relevant conjugates. Bioconj. Chem. March 1995, Vol. 6, No. 2, pages 150-165.	1-4, 11-13, 21, 28, 37, 49-51, 62 and 63
A	INADA et al. Biomedical and biotechnological applications of PEG- and PM-modified proteins. Trends Biotechnol. March 1995, Vol. 13, No. 3, pages 86-91.	1-4, 11-13, 21, 28, 37, 49-51, 62 and 63

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/08654

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
1-4, 11-13, 21, 28, 37, 49-51, 62-63
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☒

- The additional search fees were accompanied by the applicant's protest.
No protest accompanied the payment of additional search fees.

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

424/193.1, 178.1; 435/7.1, 69.6, 69.7, 69.1; 530/387.1, 810, 807

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-4, 11-13, 21, 28 and 37, drawn to a single chain antigen binding polypeptide conjugated to a polyalkylene oxide (including the one wherein specific cysteine residues have been modified), a method of producing the same and a method of use in detecting an antigen.

Group II, claim(s) 1-4, 49-51, 62 and 63, drawn to a single chain antigen binding polypeptide conjugated to a polyalkylene oxide (including the one wherein specific lysine residues have been modified).

Group III, claim(s) 5-10, 22-27 and 29-36, drawn to a single chain antigen binding polypeptide capable of polyalkylene oxide conjugation and a method of producing the same.

Group IV, claim(s) 14-20, drawn to a polynucleotide sequence, an expression vehicle containing the sequence, a host cell containing the said expression vehicle wherein the polynucleotide sequence encodes a single chain antigen binding polypeptide capable of polyalkylene oxide conjugation.

Group V, claim(s) 38 and 39, drawn to a method of imaging employing polyalkylene oxide conjugated single chain antigen binding polypeptide.

Group VI, claim 40, drawn to a method of treating a disease by administering the polyalkylene oxide conjugated single chain antigen binding polypeptide.

Group VII, claim(s) 41, 48, 52-56 and 57-61, drawn to a single chain antigen binding polypeptide capable of polyalkylene oxide conjugation wherein plural lysines are employed as sites for conjugation and a method of making the same.

Group VIII, claim(s) 42-47, drawn to a polynucleotide sequence encoding a single chain antigen binding polypeptide capable of polyalkylene oxide conjugation wherein plural lysines are employed as sites for conjugation,

The inventions listed as Groups I-VIII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The expression "special technical features" shall mean those technical features that define a contribution which each of the claimed inventions, considered as a whole, makes over the prior art.

The unifying feature of Group I or Group II appears to be the conjugate comprised of a single chain antigen binding polypeptide conjugated with polyalkylene oxide. Such conjugates are suggested by the combined teachings of Zalipsky (US 5,122,614) which disclose PEG conjugation of polypeptides for increasing their stability and Ladner et al. (US 4,946,778) which disclose single chain antigen binding polypeptides. One of ordinary skill in the art would have found it obvious to have conjugated single chain antigen binding polypeptides with PEG as described by Zalipsky so as to increase their stability. Since the apparent unifying feature is suggested in the prior art, unity of invention does not exist between Groups I to VIII, because the unifying feature is not a special technical feature.

It is further noted that technically the absence of a special technical feature would permit the separation of the conjugate of Group I from its method of making (claim 21) and a method for its use (claim 37).

Group IV, claims 14-20, appears to be suggested by the teachings of Ladner et al. which are directed to the recombinant expression of a single chain antigen binding polypeptide using an expression vehicle and a host cell.

In defining each of the groups the apparent unifying feature was set forth and it is clear that the unifying feature of each of Groups II, III and V-VIII is different and distinct from each other.